

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
12 September 2002 (12.09.2002)

PCT

(10) International Publication Number  
**WO 02/070029 A2**

(51) International Patent Classification<sup>7</sup>: **A61L 27/12,**  
27/56

(21) International Application Number: PCT/US02/05827

(22) International Filing Date: 26 February 2002 (26.02.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/798,518 2 March 2001 (02.03.2001) US  
09/960,789 21 September 2001 (21.09.2001) US

(71) Applicant (*for all designated States except US*):  
**STRYKER CORPORATION** [US/US]; 2725 Fair-  
field Road, Kalamazoo, MI 49005 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **DALAL, Paresh,**  
S. [IN/US]; 465 Boston Turnpike, F-4, Shrewsbury, MA  
01545 (US). **DIMAANO, Godofredo, R.** [US/US]; 87

Brookhill Avenue, Edison, NJ 08817 (US). **TOTH, Carol,**  
Ann [US/US]; 24 Upland Road, Sharon, MA 02067 (US).  
**KULKARNI, Shailesh, C.** [IN/US]; 6 Post Oak Lane,  
Apartment 12, Natick, MA 01760 (US).

(74) Agents: **HALEY, James, F.** et al.; c/o Fish & Neave, 1251  
Avenue of the Americas, New York, NY 10020 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

[Continued on next page]

(54) Title: POROUS BETA-TRICALCIUM PHOSPHATE GRANULES AND METHODS FOR PRODUCING SAME



(57) Abstract: A porous  $\beta$ -tricalcium phosphate material for bone implantation is provided. The multiple pores in the porous TCP body are separate discrete voids and are not interconnected. The pore size diameter is in the range of 20-500  $\mu$ m, preferably 50-125  $\mu$ m. The porous  $\beta$ -TCP material provides a carrier matrix for bioactive agents and can form a moldable putty composition upon the addition of a binder. Preferably, the bioactive agent is encapsulated in a biodegradable agent. The invention provides a kit and an implant device comprising the porous  $\beta$ -TCP, and a bioactive agent and a binder. The invention also provides an implementable prosthetic device comprising a prosthetic implant having a surface region, a porous  $\beta$ -TCP material disposed on the surface region optionally comprising at least a bioactive agent or a binder. Methods of producing the porous  $\beta$ -TCP material and including bone formation are also provided.

WO 02/070029 A2

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

**Published:**

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**Porous beta-tricalcium phosphate granules  
and methods for producing same**

BACKGROUND OF THE INVENTION

[0001] Bone tissue in the human body comprises the  
5 largest proportion of the body's connective tissue mass.  
However, unlike other connective tissues, its matrix  
consists of physiologically mineralized, tiny  
crystallites of a basic, carbonate-containing calcium  
phosphate called hydroxyapatite distributed in an  
10 organized collagen structure. Repair of this tissue is a  
complex process involving a number of cellular functions  
directed towards the formation of a scaffold and  
mineralization of the defect followed by an eventual  
remodeling of the defect site to attain the original  
15 structure.

[0002] Implantations of calcium phosphate based  
biomaterials have been found to be generally compatible  
and conducive to bone repair. Bone repair is influenced  
by a number of physico-chemical variables associated with  
20 calcium phosphate such as the calcium to phosphate molar  
ratio. Hydroxyapatite and tricalcium phosphate are  
widely used in bone implants. Hydroxyapatite has the  
chemical formula  $\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$ , and the ratio of calcium  
to phosphate is about 1.67. Tricalcium phosphate (TCP)

- 2 -

has the formula of  $\text{Ca}_3(\text{PO}_4)_2$ , and the ratio of calcium to phosphate is about 1.5. Tricalcium phosphate has biological properties of being non-reactive and resorbable. It acts as a scaffolding for bone ingrowth and undergoes progressive degradation and replacement by bone (Lange et al., Annals of Clinical and Laboratory Science, 16, pp. 467-472 (1986)). TCP is degraded 10-20 times faster than hydroxyapatite. A TCP implant generally results in superior remodeling than hydroxyapatite during the final stage of bone formation. It is noteworthy that TCP is resorbed by osteoclast cells, whereas, the much slower resorption of hydroxyapatite is effected mainly by foreign-body giant cells. The giant cells have a limit as to the amount of hydroxyapatite they will resorb.

[0003] Porous ceramic material is often selected as the matrix for bone implants. When such material is embedded at the implant site, the porous material is resorbed by osteolytic cells which infiltrate the pores. Simultaneously, the bone tissue is regenerated by osteoblasts. A certain pore size is required for osteoblasts to invade the pore of the implant material. Parameters such as crystallinity, solubility, particle size, porosity, pore structure and pore size of the implanted material can greatly influence bone compatibility and bone integration. An inappropriate combination of the above parameters can lead to improper bone repair.

[0004] The use of porous ceramics having interconnected pores as an implantable solid material for bone substitutes has been described (see, e.g., U.S. patent 5,171,720; see also Frayssinet et al., Biomaterials, 14, pp. 423-429 (1993)). Such porous ceramics, however, are brittle and are not capable of



- 3 -

being easily shaped by the practitioner during an operation.

5 [0005] Excessively large pore size and high porosity of the ceramic material can lead to excessive resorption rates, thus, preventing the matrix from providing a scaffold for the newly synthesized bone. When the rate of resorption is faster than the rate of bone growth, it often leads to an inflammatory response. Small pore size and low porosity of the ceramic material will lead to low  
10 resorption rates causing encapsulation of matrix particles in the new bone.

[0006] It would thus be desirable to identify a biomaterial which can be applied to a defect site and which can greatly enhance the regenerative process,  
15 particularly when used with other bioactive agents such as bone morphogenic proteins and other related factors. In addition, it would be desirable to identify and use a matrix which acts as a mechanically durable carrier for the bioactive agents and is a well-tolerated bone  
20 replacement material that favors healing.

#### SUMMARY OF THE INVENTION

[0007] The present invention solves these problems by identifying a porous ceramic material having a composition, pore size, porosity and granule size for  
25 improving the regeneration of bone tissue in a living body, and repairing a bone defect in a human or animal. The present invention provides a porous  $\beta$ -tricalcium phosphate ( $\beta$ -TCP) material for use in bone implant applications. The invention provides porous forms of  $\beta$ -  
30 TCP granules which are biocompatible and support the development of new bone throughout its structural form.

[0008] The invention also provides a composition comprising the porous  $\beta$ -TCP with a bioactive agent such

- 4 -

as an antibiotic, a bone morphogenic protein (BMP), or a nucleic acid molecule comprising a sequence encoding BMP in the presence or absence of a morphogenic protein stimulatory factor (MPSF) to improve osteoconductivity.

5 In a preferred embodiment, the bioactive agent is encapsulated in a biodegradable agent. Preferably, the particle size of the biodegradable agent is 20-500  $\mu\text{m}$ . The porous  $\beta$ -TCP material or porous  $\beta$ -TCP/bioactive agent mixture can also be used in conjunction with binders to  
10 form a moldable putty composition ready for shaping in the implant site. The invention also provides a kit comprising the porous  $\beta$ -TCP, and at least one or more additional components including a bioactive agent and a binder.

15 [0009] In another aspect, the invention also provides an implantable device comprising the porous  $\beta$ -TCP material, and optionally comprising one or more additional components including a bioactive agent such as a BMP, an antibiotic or a binder. The invention also  
20 provides an implantable prosthetic device comprising the porous  $\beta$ -TCP material and optionally comprising one or more additional components including a bioactive agent such as a BMP, an antibiotic or a binder. The prosthetic device or implantable device comprising the porous  $\beta$ -TCP  
25 and BMP may optionally comprise a MPSF.

[0010] Another object of the invention is to provide a method of producing the porous  $\beta$ -TCP material. The method comprises blending the TCP powder with a pore-forming agent, adding a granulating solution to form a  
30 crumbly mass, passing the crumbly mass through a sieve to form granules and sintering the granules to form the porous  $\beta$ -TCP.

[0011] The invention also provides a method of inducing bone formation in a mammal comprising the step

- 5 -

of implanting in the defect site of a mammal a composition comprising the porous  $\beta$ -TCP and optionally a binder and/or a bioactive agent. The invention describes a method of delivering a bioactive agent at a site requiring bone formation comprising implanting at the defect site of a mammal a composition comprising the porous  $\beta$ -TCP and a bioactive agent, wherein the bioactive agent is optionally encapsulated in a biodegradable agent. The invention also describes a method of delivering a bioactive agent to a site requiring cartilage formation comprising implanting at the defect site of a mammal a composition comprising the bioactive agent and biodegradable agent, wherein the bioactive agent is encapsulated in the biodegradable agent.

BRIEF DESCRIPTION OF THE DRAWINGS

[0012] Figure 1. Histologic image of animal number 297L (left tibia) at 4 weeks with placebo. From top to bottom, the sites are proximal, middle and distal, each containing  $\beta$ -TCP putty 89B,  $\beta$ -TCP putty 89C,  $\beta$ -TCP putty 89F, respectively.

[0013] Figure 2. Histologic image of animal number 297R (right tibia) at 4 weeks with placebo. From top to bottom, the sites are proximal, middle and distal, each containing control, collagen 48C,  $\beta$ -TCP putty 89A, respectively.

[0014] Figure 3. Histologic image of animal number 295L (left tibia) at 4 weeks with OP-1. From top to bottom, the sites are proximal, middle and distal, each containing collagen 48C,  $\beta$ -TCP putty 89A,  $\beta$ -TCP putty 89B, respectively.

[0015] Figure 4. Histologic image of animal number 295R (right tibia) at 4 weeks with OP-1. From top to bottom, the sites are proximal, middle and distal, each

- 6 -

containing  $\beta$ -TCP putty 89C,  $\beta$ -TCP putty 89F, control, respectively.

5 [0016] Figure 5. Histologic image of animal number 299L (left tibia) at 8 weeks with placebo. From top to bottom, the sites are proximal, middle and distal, each containing  $\beta$ -TCP putty 89B,  $\beta$ -TCP putty 89C,  $\beta$ -TCP putty 89F, respectively.

10 [0017] Figure 6. Histologic image of animal number 299R (right tibia) at 8 weeks with placebo. From top to bottom, the sites are proximal, middle and distal, each containing control, collagen 48C,  $\beta$ -TCP putty 89A, respectively.

15 [0018] Figure 7. Histologic image of animal number 138L (left tibia) at 8 weeks with OP-1. From top to bottom, the sites are proximal, middle and distal, each containing  $\beta$ -TCP putty 89A,  $\beta$ -TCP putty 89B,  $\beta$ -TCP putty 89C, respectively.

20 [0019] Figure 8. Histologic image of animal number 138R (right tibia) at 8 weeks with OP-1. From top to bottom, the sites are proximal, middle and distal, each containing  $\beta$ -TCP putty 89F, control, collagen 48C, respectively.

25 [0020] Figure 9. Radiographic image of animal number 297L (left tibia) at 4 weeks with placebo. From the right, the sites are proximal, middle and distal, each containing  $\beta$ -TCP putty 89B,  $\beta$ -TCP putty 89C,  $\beta$ -TCP putty 89F, respectively.

30 [0021] Figure 10. Radiographic image of animal number 297R (right tibia) at 4 weeks with placebo. From the left, the sites are proximal, middle and distal, each containing control, collagen 48C,  $\beta$ -TCP putty 89A, respectively.

[0022] Figure 11. Radiographic image of animal number 295L (left tibia) at 4 weeks with OP-1. From the left,

- 7 -

the sites are proximal, middle and distal, each containing collagen 48C,  $\beta$ -TCP putty 89A,  $\beta$ -TCP putty 89B, respectively.

5 [0023] Figure 12. Radiographic image of animal number 295R (right tibia) at 4 weeks with OP-1. From the left, the sites are proximal, middle and distal, each containing  $\beta$ -TCP putty 89C,  $\beta$ -TCP putty 89F, control, respectively.

10 [0024] Figure 13. Radiographic image of animal number 299L (left tibia) at 8 weeks with placebo. From the right, the sites are proximal, middle and distal, each containing  $\beta$ -TCP putty 89B,  $\beta$ -TCP putty 89C,  $\beta$ -TCP putty 89F, respectively.

15 [0025] Figure 14. Radiographic image of animal number 299R (right tibia) at 8 weeks with placebo. From the left, the sites are proximal, middle and distal, each containing control, collagen 48C,  $\beta$ -TCP putty 89A, respectively.

20 [0026] Figure 15. Radiographic image of animal number 138L (left tibia) at 8 weeks with OP-1. From the right, the sites are proximal, middle and distal, each containing  $\beta$ -TCP putty 89A,  $\beta$ -TCP putty 89B,  $\beta$ -TCP putty 89C, respectively.

25 [0027] Figure 16. Radiographic image of animal number 138R (right tibia) at 8 weeks with OP-1. From the left, the sites are proximal, middle and distal, each containing  $\beta$ -TCP putty 89F, control, collagen 48C, respectively.

30 [0028] Figure 17. Paraffin scanning image of animal number 297L (left tibia) at 4 weeks with placebo. From the top, the sites are proximal, middle and distal, each containing  $\beta$ -TCP putty 89B,  $\beta$ -TCP putty 89C,  $\beta$ -TCP putty 89F, respectively.

- 8 -

[0029] Figure 18. Paraffin scanning image of animal number 297R (right tibia) at 4 weeks with placebo. From the top, the sites are proximal, middle and distal, each containing control, collagen 48C,  $\beta$ -TCP putty 89A, respectively.

[0030] Figure 19. Paraffin scanning image of animal number 295L (left tibia) at 4 weeks with OP-1. From the top, the sites are proximal, middle and distal, each containing collagen 48C,  $\beta$ -TCP putty 89A,  $\beta$ -TCP putty 89B, respectively.

[0031] Figure 20. Paraffin scanning image of animal number 295R (right tibia) at 4 weeks with OP-1. From the top, the sites are proximal, middle and distal, each containing  $\beta$ -TCP putty 89C,  $\beta$ -TCP putty 89F, control, respectively.

[0032] Figure 21. Paraffin scanning image of animal number 299L (left tibia) at 8 weeks with placebo. From the top, the sites are proximal, middle and distal, each containing  $\beta$ -TCP putty 89B,  $\beta$ -TCP putty 89C,  $\beta$ -TCP putty 89F, respectively.

[0033] Figure 22. Paraffin scanning image of animal number 299R (right tibia) at 8 weeks with placebo. From the top, the sites are middle and distal, each containing collagen 48C and  $\beta$ -TCP putty 89A, respectively.

[0034] Figure 23. Paraffin scanning image of animal number 138L (left tibia) at 8 weeks with OP-1. From the top, the sites are proximal, middle and distal, each containing  $\beta$ -TCP putty 89A,  $\beta$ -TCP putty 89B,  $\beta$ -TCP putty 89C, respectively.

[0035] Figure 24. Paraffin scanning image of animal number 138R (right tibia) at 8 weeks with OP-1. From the top, the sites are proximal, middle and distal, each containing  $\beta$ -TCP putty 89F, control, collagen 48C, respectively.

- 9 -

[0036] Figure 25. Specimen 295L middle site showing one of the five pores with bone growth, where EP is an empty pore and FP is a filled pore.

5 [0037] Figure 26. Specimen 299L distal site showing 7 or 8 pores with bone growth, where EP is any empty pore and FP is a filled pore.

10 [0038] Figure 27. Radiographic image of animal number 5333L (left tibia) at 4 weeks with OP-1 encapsulated in PLGA. From the left, the sites are proximal, middle and distal, each containing control, formulation 5 and formulation 4, respectively.

15 [0039] Figure 28. Radiographic image of animal number 5335L (left tibia) at 8 weeks with OP-1 encapsulated in PLGA. From the left, the sites are proximal, middle and distal, each containing control, formulation 4 and formulation 5, respectively.

#### DETAILED DESCRIPTION OF THE INVENTION

20 [0040] In order that the invention herein described may be fully understood, the following detailed description is set forth.

25 [0041] "Amino acid sequence homology" is understood to include both amino acid sequence identity and similarity. Homologous sequences share identical and/or similar amino acid residues, where similar residues are conservative substitutions for, or "allowed point mutations" of, corresponding amino acid residues in an aligned reference sequence. Thus, a candidate polypeptide sequence that shares 70% amino acid homology with a reference sequence is one in which any 70% of the aligned residues are  
30 either identical to, or are conservative substitutions of, the corresponding residues in a reference sequence. Certain particularly preferred morphogenic polypeptides share at least 60%, and preferably 70% amino acid

- 10 -

sequence identity with the C-terminal 102-106 amino acids, defining the conserved seven-cysteine domain of human OP-1, BMP-2, and related proteins.

[0042] Amino acid sequence homology can be determined  
5 by methods well known in the art. For instance, to determine the percent homology of a candidate amino acid sequence to the sequence of the seven-cysteine domain, the two sequences are first aligned. The alignment can be made with, e.g., the dynamic programming algorithm  
10 described in Needleman et al., J. Mol. Biol., 48, pp. 443 (1970), and the Align Program, a commercial software package produced by DNASTar, Inc. The teachings by both sources are incorporated by reference herein. An initial alignment can be refined by comparison to a multi-  
15 sequence alignment of a family of related proteins. Once the alignment is made and refined, a percent homology score is calculated. The aligned amino acid residues of the two sequences are compared sequentially for their similarity to each other. Similarity factors include  
20 similar size, shape and electrical charge. One particularly preferred method of determining amino acid similarities is the PAM250 matrix described in Dayhoff et al., Atlas of Protein Sequence and Structure, 5, pp. 345-352 (1978 & Supp.), which is incorporated herein by  
25 reference. A similarity score is first calculated as the sum of the aligned pairwise amino acid similarity scores. Insertions and deletions are ignored for the purposes of percent homology and identity. Accordingly, gap penalties are not used in this calculation. The raw  
30 score is then normalized by dividing it by the geometric mean of the scores of the candidate sequence and the seven-cysteine domain. The geometric mean is the square root of the product of these scores. The normalized raw score is the percent homology.



- 11 -

[0043] "Biocompatible" refers to a material that does not elicit detrimental effects associated with the body's various protective systems, such as cell and humoral-associated immune responses, e.g., inflammatory responses and foreign body fibrotic responses. The term  
5 biocompatible also implies that no specific undesirable cytotoxic or systemic effects are caused by the material when it is implanted into the patient.

[0044] "Binder" refers to any biocompatible material  
10 which, when admixed with osteogenic protein and/or the porous matrix promotes bone formation. Certain preferred binders promote such repair using less osteogenic protein than standard osteogenic devices. Other preferred binders can promote repair using the same amount of the  
15 osteogenic protein as the standard osteogenic devices while some require more to promote repair. As taught herein, the skilled artisan can determine an effective amount of protein for use with any suitable binder using only routine experimentation. Among the other  
20 characteristics of a preferred binder is an ability to render the device: pliable, shapeable and/or malleable; injectable; adherent to bone, cartilage, muscle and other tissues, resistant to disintegration upon washing and/or irrigating during surgery; and, resistant to dislodging  
25 during surgery, suturing and post-operatively, to name but a few. Additionally, in certain preferred embodiments, a binder can achieve the aforementioned features and benefits when present in low proportions.

[0045] "Biodegradable agent" refers to a resorbable  
30 biocompatible material such as a material that degrades gradually at the implant site. The material is capable of encapsulating a bioactive agent to provide time release or sustained release delivery of the bioactive agent. The biodegradable material encompasses natural

- 12 -

and synthetic polymers. Examples of biodegradable material are poly(L-lactide) (PLLA), poly(D,L-lactide) (PDLA), polyglycolide (PGA), poly(lactide-co-glycolide (PLGA) and co-polymers thereof.

5 [0046] "Bone" refers to a calcified (mineralized) connective tissue primarily comprising a composite of deposited calcium and phosphate in the form of hydroxyapatite, collagen (primarily Type I collagen) and bone cells such as osteoblasts, osteocytes and  
10 osteoclasts, as well as to bone marrow tissue which forms in the interior of true endochondral bone. Bone tissue differs significantly from other tissues, including cartilage tissue. Specifically, bone tissue is vascularized tissue composed of cells and a biphasic  
15 medium comprising a mineralized, inorganic component (primarily hydroxyapatite crystals) and an organic component (primarily of Type I collagen). Glycosaminoglycans constitute less than 2% of this organic component and less than 1% of the biphasic medium  
20 itself, or of bone tissue per se. Moreover, relative to cartilage tissue, the collagen present in bone tissue exists in a highly-organized parallel arrangement. Bony defects, whether from degenerative, traumatic or cancerous etiologies, pose a formidable challenge to the  
25 reconstructive surgeon. Particularly difficult is reconstruction or repair of skeletal parts that comprise part of a multi-tissue complex, such as occurs in mammalian joints.

[0047] "Bone formation" means formation of  
30 endochondral bone or formation of intramembranous bone. In humans, bone formation begins during the first 6-8 weeks of fetal development. Progenitor stem cells of mesenchymal origin migrate to predetermined sites, where they either: (a) condense, proliferate, and

- 13 -

differentiate into bone-forming cells (osteoblasts), a process observed in the skull and referred to as "intramembranous bone formation" or, (b) condense, proliferate and differentiate into cartilage-forming cells (chondroblasts) as intermediates, which are subsequently replaced with bone-forming cells. More specifically, mesenchymal stem cells differentiate into chondrocytes. The chondrocytes then become calcified, undergo hypertrophy and are replaced by newly formed bone made by differentiated osteoblasts, which now are present at the site. Subsequently, the mineralized bone is extensively remodeled, thereafter becoming occupied by an ossicle filled with functional bone-marrow elements. This process is observed in long bones and referred to as "endochondral bone formation." In postfetal life, bone has the capacity to repair itself upon injury by mimicking the cellular process of embryonic endochondral bone development. That is, mesenchymal progenitor stem cells from the bone-marrow, periosteum, and muscle can be induced to migrate to the defect site and begin the cascade of events described above. There, they accumulate, proliferate, and differentiate into cartilage, which is subsequently replaced with newly formed bone.

[0048] "Bone morphogenic protein (BMP)" refers to a protein belonging to the BMP family of the TGF- $\beta$  superfamily of proteins (BMP family) based on DNA and amino acid sequence homology. A protein belongs to the BMP family according to this invention when it has at least 50% amino acid sequence identity with at least one known BMP family member within the conserved C-terminal cysteine-rich domain which characterizes the BMP protein family. Members of the BMP family may have less than 50% DNA or amino acid sequence identity overall.

- 14 -

[0049] "Conservative substitutions" are residues that are physically or functionally similar to the corresponding reference residues. That is, a conservative substitution and its reference residue have similar size, shape, electric charge, chemical properties including the ability to form covalent or hydrogen bonds, or the like. Preferred conservative substitutions are those fulfilling the criteria defined for an accepted point mutation in Dayhoff et al., supra. Examples of conservative substitutions are substitutions within the following groups: (a) valine, glycine; (b) glycine, alanine; (c) valine, isoleucine, leucine; (d) aspartic acid, glutamic acid; (e) asparagine, glutamine; (f) serine, threonine; (g) lysine, arginine, methionine; and (h) phenylalanine, tyrosine. The term "conservative variant" or "conservative variation" also includes the use of a substituting amino acid residue in place of an amino acid residue in a given parent amino acid sequence, where antibodies specific for the parent sequence are also specific for, i.e., "cross-react" or "immuno-react" with, the resulting substituted polypeptide sequence.

[0050] "Defect" or "defect site" refers to a site requiring bone, joint, cartilage or ligament repair, construction, fusion, regeneration or augmentation. The site may be an orthopedic structural disruption or abnormality, or a site where bone does not normally grow. The defect further can define an osteochondral defect, including a structural disruption of both the bone and overlying cartilage. A defect can assume the configuration of a "void", which is understood to mean a three-dimensional defect such as, for example, a gap, cavity, hole or other substantial disruption in the structural integrity of a bone or joint. A defect can be the result of accident, disease, surgical manipulation,

- 15 -

and/or prosthetic failure. In certain embodiments, the defect is a void having a volume incapable of endogenous or spontaneous repair. Such defects in long bone are generally twice the diameter of the subject bone and are also called "critical size" defects. For example, in a canine ulna defect model, the art recognizes such defects to be approximately 3-4 cm. Generally, critical size defects are approximately 1.0 cm, and incapable of spontaneous repair. See, for example, Schmitz et al., Clinical Orthopaedics and Related Research, 205, pp. 299-308 (1986); and Vukicevic et al., in Advances in Molecular and Cell Biology, 6, pp. 207-224 (1993) (JAI Press, Inc.). In rabbit and monkey segmental defect models, the gap is approximately 1.5 cm and 2.0 cm, respectively. In other embodiments, the defect is a non-critical size segmental defect. Generally, these are capable of spontaneous repair. In certain other embodiments, the defect is an osteochondral defect, such as an osteochondral plug. Such a defect traverses the entirety of the overlying cartilage and enters, at least in part, the underlying bony structure. In contrast, a chondral or subchondral defect traverses the overlying cartilage, in part or in whole, respectively, but does not involve the underlying bone. Other defects susceptible to repair using the instant invention include, but are not limited to, non-union fractures; bone cavities; tumor resection; fresh fractures (distracted or undistracted); cranial, maxillofacial and facial abnormalities, for example, in facial skeletal reconstruction, specifically, orbital floor reconstruction, augmentation of the alveolar ridge or sinus, periodontal defects and tooth extraction socket; cranioplasty, genioplasty, chin augmentation, palate reconstruction, and other large bony reconstructions;

- 16 -

vertebroplasty, interbody fusions in the cervical, thoracic and lumbar spine and posteriolateral fusions in the thoracic and lumbar spine; in osteomyelitis for bone regeneration; appendicular fusion, ankle fusion, total  
5 hip, knee and joint fusions or arthroplasty; correcting tendon and/or ligamentous tissue defects such as, for example, the anterior, posterior, lateral and medial ligaments of the knee, the patella and achilles tendons, and the like as well as those defects resulting from  
10 diseases such as cancer, arthritis, including osteoarthritis, and other bone degenerative disorders such as osteochondritis dessicans.

[0051] "Granulating solution" refers to a solution that has a certain degree of consistency and  
15 cohesiveness, and enhances the formation of granules.

[0052] "Morphogenic protein" refers to a protein having morphogenic activity (see below). Preferably a morphogenic protein of this invention comprises at least one polypeptide belonging to the BMP protein family.  
20 Morphogenic proteins may be capable of inducing progenitor cells to proliferate and/or to initiate differentiation pathways that lead to cartilage, bone, tendon, ligament, neural or other types of tissue formation depending on local environmental cues, and thus  
25 morphogenic proteins may behave differently in different surroundings. For example, an osteogenic protein may induce bone tissue at one treatment site and neural tissue at a different treatment site.

[0053] "Morphogenic protein stimulatory factor (MPSF)"  
30 refers to a factor that is capable of stimulating the ability of a morphogenic protein to induce tissue formation from a progenitor cell. The MPSF may have a direct or indirect effect on enhancing morphogenic protein inducing activity. For example, the MPSF may

- 17 -

increase the bioactivity of another MPSF. Agents that increase MPSF bioactivity include, for example, those that increase the synthesis, half-life, reactivity with other biomolecules such as binding proteins and receptors, or the bioavailability of the MPSF.

[0054] "Osteogenic protein (OP)" refers to a morphogenic protein that is capable of inducing a progenitor cell to form cartilage and/or bone. The bone may be intramembranous bone or endochondral bone. Most osteogenic proteins are members of the BMP protein family and are thus also BMPs. As described elsewhere herein, the class of proteins is typified by human osteogenic protein (hOP-1). Other osteogenic proteins useful in the practice of the invention include osteogenically active forms of OP-1, OP-2, OP-3, BMP-2, BMP-3, BMP-4, BMP-5, BMP-6, BMP-9, DPP, Vgl, Vgr, 60A protein, GDF-1, GDF-3, GDF-5, GDF-6, GDF-7, BMP-10, BMP-11, BMP-13, BMP-15, UNIVIN, NODAL, SCREW, ADMP or NEURAL and amino acid sequence variants thereof. In one currently preferred embodiment, osteogenic protein includes any one of: OP-1, OP-2, OP-3, BMP-2, BMP-4, BMP-5, BMP-6, BMP-9, and amino acid sequence variants and homologs thereof, including species homologs thereof. Particularly preferred osteogenic proteins are those comprising an amino acid sequence having at least 70% homology with the C-terminal 102-106 amino acids, defining the conserved seven cysteine domain, of human OP-1, BMP-2, and related proteins. Certain preferred embodiments of the instant invention comprise the osteogenic protein, OP-1. As further described elsewhere herein, the osteogenic proteins suitable for use with applicants' invention can be identified by means of routine experimentation using the art-recognized bioassay described by Reddi and

- 18 -

Sampath (Sampath et al., Proc. Natl. Acad. Sci., 84, pp. 7109-13, incorporated herein by reference)

[0055] Proteins useful in this invention include eukaryotic proteins identified as osteogenic proteins (see U.S. Patent 5,011,691, incorporated herein by reference), such as the OP-1, OP-2, OP-3 and CBMP-2 proteins, as well as amino acid sequence-related proteins, such as DPP (from *Drosophila*), Vg1 (from *Xenopus*), Vgr-1 (from mouse), GDF-1 (from humans, see Lee, PNAS, 88, pp. 4250-4254 (1991)), 60A (from *Drosophila*, see Wharton et al. PNAS, 88, pp. 9214-9218 (1991)), dorsalin-1 (from chick, see Basler et al. Cell 73, pp. 687-702 (1993) and GenBank accession number L12032) and GDF-5 (from mouse, see Storm et al. Nature, 368, pp. 639-643 (1994)). The teachings of the above references are incorporated herein by reference. BMP-3 is also preferred. Additional useful proteins include biosynthetic morphogenic constructs disclosed in U.S. Pat. No. 5,011,691, incorporated herein by reference, e.g., COP-1, COP-3, COP-4, COP-5, COP-7 and COP-16, as well as other proteins known in the art. Still other proteins include osteogenically active forms of BMP-3b (see Takao, et al. Biochem. Biophys. Res. Comm., 219, pp. 656-662 (1996)). BMP-9 (see WO95/33830), BMP-15 (see WO96/35710), BMP-12 (see WO95/16035), CDMP-1 (see WO 94/12814), CDMP-2 (see WO94/12814), BMP-10 (see WO94/26893), GDF-1 (see WO92/00382), GDF-10 (see WO95/10539), GDF-3 (see WO94/15965) and GDF-7 (see WO95/01802). The teachings of the above references are incorporated herein by reference.

[0056] "Repair" is intended to mean new bone and/or cartilage formation which is sufficient to at least partially fill the void or structural discontinuity at the defect. Repair does not, however, mean, or otherwise



- 19 -

necessitate, a process of complete healing or a treatment which is 100% effective at restoring a defect to its pre-defect physiological/structural/mechanical state.

[0057] "Synergistic interaction" refers to an  
5 interaction in which the combined effect of two or more agents is greater than the algebraic sum of their individual effects.

#### Porous $\beta$ -TCP

10 [0058] This present invention provides a porous  $\beta$ -TCP having a pore size and granule size appropriate for bone formation, bone regeneration, and bone repair at a defect site in a human or animal. The porous  $\beta$ -TCP body  
described in this invention comprises  $\beta$ -TCP having a  
15 multiplicity of pores. Each pore is a single separate void partitioned by walls and is not interconnected. The porous  $\beta$ -TCP body of this invention is distinct from the cancellous or fenestrate structures that contain  
capillary void paths or interconnections between adjacent  
20 pores. The pore diameter size of the porous  $\beta$ -TCP of this invention is in the range of 20-500  $\mu\text{m}$ . In one embodiment, the pore diameter size is in the range of 410-460  $\mu\text{m}$ . In a preferred embodiment, the pore diameter size is 40-190  $\mu\text{m}$ . In another embodiment, the pore  
25 diameter size is in the range of 20-95  $\mu\text{m}$ . In a more preferred embodiment, the pore diameter is in the range of 50-125  $\mu\text{m}$ . These pores provide residence spaces for the infiltrating osteolytic cells and osteoblasts when the  $\beta$ -TCP material is embedded in the living body. In  
30 one embodiment, the pores are spherical and uniformly distributed. Spherical pores having a diameter in the range of 20-500  $\mu\text{m}$  are appropriate for osteoblast infiltration. Spherical pores also provide the porous body with the necessary mechanical strength during the

- 20 -

period that new bone is being synthesized, thus preventing the bone from fracturing during this period.

[0059] Tricalcium phosphate (TCP) has the formula of  $\text{Ca}_3(\text{PO}_4)_2$ , with the Ca/P ratio being about 1.5. TCP powder has an apatite crystal structure. Upon sintering, the apatite structure converts to the rhombic  $\beta$ -TCP structure. At high temperatures, the metastable,  $\alpha$ -TCP structure can also form.  $\alpha$ -TCP is known to have excessive solubility, which does not permit the rate of resorption to be complementary to the rate of substitution by the hard tissue. In addition,  $\alpha$ -TCP is capable of generating harmful inflammatory responses. In a preferred embodiment, the TCP is sintered at high temperatures of 1100-1200°C. Above 1300°C, TCP is converted to the metastable  $\alpha$ -TCP. Sintering the TCP reduces its solubility in body fluids, which leads to a corresponding reduction in its chemical activity so that the porous TCP is well tolerated in the body and acute inflammatory reactions are avoided. Therefore, the porous  $\beta$ -TCP is preferably sintered. More preferably the  $\beta$ -TCP comprises  $\beta$ -TCP that is 95-100% pure.

[0060] The porous  $\beta$ -TCP material of the present invention may have any shape and size. In one embodiment, the porous  $\beta$ -TCP is granular and has a particle size between 0.1 to 2 mm. In a preferred embodiment, the particle size is 0.5-1.7 mm. In a more preferred embodiment the particle size is 1.0-1.7 mm. In a most preferred embodiment, the particle size is 0.5-1 mm.  $\beta$ -TCP having a granule size of less than 0.1 mm is not appropriate because it will be readily displaced by flowing body fluids. On the other hand, although bone formation is more obvious in larger particles,  $\beta$ -TCP having a granule size greater than 2 mm is also not appropriate because too many or excessively large gaps

- 21 -

will form between the granules, thus preventing the effective coalescence of the  $\beta$ -TCP to the newly synthesized bone.

5 [0061] The porosity of the  $\beta$ -TCP influences the resorption rate. If the porosity is too high, the strength of the granules will be decreased. If the porosity is too low, the rate of resorption will be slow. The total porosity is measured using the mercury intrusion parameter method or equivalent methods. In one  
10 embodiment, the total porosity is in the range of 5-80%. In another embodiment, the total porosity is in the range of 40-80%. In a more preferred embodiment, the total porosity is 65-75%. In a most preferred embodiment, the total porosity is 70%.

15 [0062] The porous  $\beta$ -TCP of this invention may also be combined with one or more bioactive agents. The bioactive agent may be an agent that enhances bone growth or a substance that is medically useful or combinations thereof. It is envisioned that the bioactive agent can  
20 include but is not limited to bone morphogenic proteins, growth factors such as EGF, PDGF, IGF, FGF, TGF- $\alpha$  and TGF- $\beta$ , cytokines, MPSF, hormones, peptides, lipids, trophic agents and therapeutic compositions including antibiotics and chemotherapeutic agents, insulin,  
25 chemoattractant, chemotactic factors, enzymes, enzyme inhibitors. It is also envisioned that bioactive agents such as vitamins, cytoskeletal agents, cartilage fragments, allografts, autografts, living cells such as chondrocytes, bone marrow cells, mesenchymal stem cells,  
30 tissue transplants, immuno-suppressants may be added to the porous  $\beta$ -TCP.

[0063] In one embodiment, the bioactive agent is a bone morphogenic protein. In a preferred embodiment, the bone morphogenic protein is OP-1 (BMP-7), OP-2, OP-3,

- 22 -

COP-1, COP-3, COP-4, COP-5, COP-7, COP-16, BMP-2, BMP-3, BMP-3b, BMP-4, BMP-5, BMP-6, BMP-9, BMP-10, BMP-11, BMP-12, BMP-13, BMP-14, BMP-15, BMP-16, BMP-17, BMP-18, GDF-1, GDF-2, GDF-3, GDF-5, GDF-6, GDF-7, GDF-8, GDF-9, 5 GDF-10, GDF-11, GDF-12, MP121, dorsalin-1, DPP, Vg-1, Vgr-1, 60A protein, NODAL, UNIVIN, SCREW, ADMP, NEURAL, and TGF- $\beta$ . In a more preferred embodiment, the morphogenic protein is OP-1.

[0064] In another embodiment the morphogenic activity of the bone morphogenic protein is enhanced by the addition of a MPSF. In a preferred embodiment the MPSF is selected from the group consisting of insulin-like growth factor I (IGF-I), estradiol, fibroblast growth factor (FGF), growth hormone (GH), growth and 15 differentiation factor (GDF), hydrocortisone (HC), insulin, progesterone, parathyroid hormone (PTH), vitamin D, retinoic acid and IL-6. In a preferred embodiment, the MPSF is selected from IGF-1, IL-6, FGF, PTH. In a more preferred embodiment, the MPSF is IGF-1.

[0065] In another embodiment, the bioactive agent is preferably an antimicrobial or antibiotic including but not limited to erythromycin, bacitracin, neomycin, penicillin, polymyxin B, tetracycline, viomycin, chloromycetin and streptomycin, cefazolin, ampicillin, 25 azactam, tobramycin, clindamycin and gentamycin. The concentrations of the antibiotic to be used are well known in the art. Such antibiotics have been known and used in connection with bone cement materials. See, for example, Hoff et al., J. Bone Joint Surg., 63A, pp. 798, 30 (1981); and Dueland et al., Clin. Orthop., 169, pp. 264-268, (1982). The teachings of these two references are incorporated herein by reference.

[0066] In another preferred embodiment, the bioactive agent is a repair cell. In a preferred embodiment, the

- 23 -

repair cell is a mammalian cell, more preferably, a human cell of the same type as that of the tissue being repaired or reconstructed. Suitable examples of repair cells include bone cells such as bone marrow stem cells, osteocytes, osteoblasts, osteoclasts and bone progenitor cells. In another embodiment, the cell is transfected with a nucleic acid molecule encoding a BMP.

[0067] In yet another preferred embodiment, the bioactive agent is a nucleic acid molecule comprising a sequence encoding a BMP, preferably, OP-1 (SEQ ID NO: 10). In a preferred embodiment, the nucleic acid molecule is a RNA or DNA molecule. The nucleic acid sequence encoding the BMP may be inserted in recombinant expression vectors. Examples of vectors include but are not limited to pBR322, pH717, pH731, pH752, pH754 and pW24. SP6 vectors may be used for *in vitro* transcription of RNA. Transcription promoters useful for expressing the BMP include but are not limited to the SV40 early promoter, the adenovirus promoter (AdMLP), the mouse metallothionein-I promoter (mMT-I), the Rous sarcoma virus (RSV) long terminal repeat (LTR), the mouse mammary tumor virus long terminal repeat (MMTV-LTR), and the human cytomegalovirus major intermediate-early promoter (hCMV). The DNA sequences for all of these promoters are known in the art and are available commercially. The DNA sequence may also be inserted in the genome of a recombinant virus such as, for example recombinant adenovirus, adeno-associated virus or retrovirus. The repair cell or bone progenitor cell is then transfected or infected with the vector or virus and expresses the BMP protein. The nucleic acid sequence may transiently or stably transfect the repair cell or bone progenitor cell.

- 24 -

[0068] In one embodiment, the nucleic acid molecule is directly injected into the implant site. Preferably, the nucleic acid is trapped in a carrier selected from the group consisting of mannitol, sucrose, lactose, trehalose, liposomes, proteoliposomes that contain viral envelope proteins and polylysine-glycoprotein complexes. See, e.g., Ledley, J. Pediatrics 110, pp. 1 (1987); Nicolau et al., Proc. Natl. Acad. Sci. U.S.A., 80, pp. 1068 (1983). In another preferred embodiment, the nucleic acid is transfected or infected into target cells such as bone progenitor cells and repair cells that have been removed from the body. The transfected cell or infected cells are then re-implanted into the body.

[0069] In a most preferred embodiment, the bioactive agent is encapsulated in a biodegradable agent. As the biodegradable agent is slowly resorbed by the osteoclast cells, the encapsulated bioactive agent is gradually released into the matrix. At the implant site, one may deliver the bioactive agent through a combination of different biodegradable agents, preferably, differing in the rate of resorption, to achieve a multiple boost delivery system. In another preferred embodiment, the biodegradable agent is multi-layered. Each layer comprises a different biodegradable agent, preferably, differing in the rate of resorption. Methods of encapsulating the bioactive agent include but are not limited to the emulsion-solvent evaporation method (Grandfils et al., Journal of Biomedical Materials Research, 26, pp. 467-479 (1992)) and the method described in Herbert et al., Pharmaceutical Research, 15, pp. 357-361 (1998). The above references are incorporated herein by reference. The latter method is especially suitable for encapsulating proteins. Other methods are described in U.S. patents 6,110,503,

- 25 -

5,654,008 and 5,271,961, which are incorporated herein by reference. In a preferred embodiment, the OP-1 is stabilized by the addition of lactose during the encapsulation process.

5 [0070] The biodegradable agents of this invention may be in bead or microsphere form. The biodegradable agents can be resorbable biocompatible polymers including both natural and synthetic polymers. Natural polymers are typically absorbed by enzymatic degradation in the body,  
10 while synthetic resorbable polymers typically degrade by a hydrolytic mechanism. It is preferred that the particle size of the biodegradable agent is 20-500  $\mu\text{m}$ , preferably, 20-140  $\mu\text{m}$ , more preferably 50-140  $\mu\text{m}$ , and most preferably 75-140  $\mu\text{m}$ .

15 [0071] In one embodiment, the biodegradable agent is selected from the group consisting of ethylenevinylacetate, natural and synthetic collagen, poly(glaxanone), poly(phosphazenes), polyglactin, polyglactic acid, polyaldonic acid, polyacrylic acids,  
20 polyalkanoates, , polyorthoesters, poly(L-lactide) (PLLA), poly(D,L-lactide) (PDLA), polyglycolide (PGA), poly(lactide-co-glycolide (PLGA), poly( $\zeta$ -caprolactone), poly(trimethylene carbonate), poly(p-dioxanone), poly( $\zeta$ -caprolactone-co-glycolide),  
25 poly(glycolide-co-trimethylene carbonate) poly(D,L-lactide-co-trimethylene carbonate), polyarylates, polyhydroxybutyrate (PHB), polyanhydrides, poly(anhydride-co-imide) and co-polymers thereof, polymers of amino acids, propylene-co-fumarates, a  
30 polymer of one or more  $\alpha$ -hydroxy carboxylic acid monomers, bioactive glass compositions, admixtures thereof and any derivatives and modifications thereof. Preferably, the modification changes less than 50% of the overall structure of the polymer.

- 26 -

[0072] In a preferred embodiment, the biodegradable agent is selected from the group consisting of polyorthoesters, poly(L-lactide) (PLLA), poly(D,L-lactide) (PDLLA), polyglycolide (PGA), poly(lactide-co-glycolide) (PLGA), poly( $\zeta$ -caprolactone), poly(trimethylene carbonate), poly(p-dioxanone), poly( $\zeta$ -caprolactone-co-glycolide), poly(glycolide-co-trimethylene carbonate), poly(D,L-lactide-co-trimethylene carbonate), polyarylates and co-polymers thereof.

[0073] In another more preferred embodiment, the biodegradable agent is selected from the group consisting of poly(glaxanone), poly(phosphazenes), ethylenevinylacetate, polyglactin, polyglactic acid, polyaldonic acid, polyacrylic acids, polyalkanoates, co-polymers thereof and natural and synthetic collagen.

[0074] In yet another more preferred embodiment the biodegradable agent is selected from the group consisting of polyhydroxybutyrate (PHB), anhydrides including polyanhydrides, poly(anhydride-co-imide) and co-polymers thereof, polymers of amino acids, propylene-co-fumarates, a polymer of one or more -hydroxy carboxylic acid monomers, (e.g.  $\alpha$ -hydroxy acetic acid (glycolic acid) and/or  $\alpha$ -hydroxy propionic acid (lactic acid)), bioactive glass compositions.  $\alpha$ -hydroxy propionic acid can be employed in its d- or l- form, or as a racemic mixture.

[0075] In a most preferred embodiment the biodegradable agent is poly(lactide-co-glycolide) (PLGA). Depending upon the desired rate of release of the bioactive agent, the molar ratio of the lactide, glycolide monomers can be adjusted. In a preferred embodiment, the monomer ratio is 50:50. In general, the higher the molecular weight, the slower the biodegradation. Preferably, the molecular weight range



- 27 -

of the polymer is from about 5,000 to 500,000 daltons, more preferably 10,000 to 30,000 daltons.

Method of Producing Porous  $\beta$ -TCP

5 [0076] The invention also relates to a method of producing porous  $\beta$ -TCP granules. The TCP used in preparing the porous  $\beta$ -TCP is prepared according to known methods in the art. The TCP is harvested via a spray dryer, preferably to a particle size of less than 10  $\mu\text{m}$ .  
10 If the particle size is too large, it will interfere with the formation of pores.

[0077] The fine TCP powder is then mixed with a pore-forming agent that decomposes at high temperature into gaseous decomposition products without leaving any solid  
15 residue. The pore-forming agents of this invention may be in bead or resin form. In one embodiment, the pore-forming agents are selected from thermally decomposable material such as naphthalene, prepolymers of polyacrylates, prepolymers of polymethacrylates,  
20 polymethyl methacrylate, copolymers of methyl acrylate and methyl methacrylate and mixtures thereof, polystyrene, polyethylene glycol, crystalline cellulose powder, fibrous cellulose, polyurethanes, polyethylenes, nylon resins and acrylic resins. In a more preferred  
25 embodiment the pore-forming agent is selected from the group consisting of polymethyl methacrylate, polystyrene and polyethylene glycol. It is preferred that the pore-forming agent creates a pore size diameter of 20-500  $\mu\text{m}$ , more preferably 40-190  $\mu\text{m}$ , and most preferably 50-125  $\mu\text{m}$   
30 after sintering.

[0078] The proportion and particle size of the pore-forming agent influences the porosity and the pore structure. An excessive amount of the pore-forming agent leads to interconnected pores and a decrease in density

- 28 -

of the  $\beta$ -TCP body and hence mechanical strength of the sintered body. A deficiency in the amount of the pore-forming agent may result in the insufficiently developed pore structure. The proportion of pore-forming agent is preferably 10-50% by weight, more preferably 30-40% by weight, most preferably 37.5% by weight.

[0079] A granulating solution is then added to the mixture of TCP powder and pore-forming agent to produce a crumbly mass. This improves the sieving procedure that follows. Depending on the desired viscosity to be achieved and the aqueous properties of the dispersing medium, the compound used to form the granulating solution may be selected from the group consisting of polyvinyl pyrrolidone, starch, gelatin, polyvinyl alcohol, polyethylene oxide, hydroxyethyl cellulose, polyvinyl butyral and cellulose acetate butyrate. Preferably, the compound in the granulating solution is selected from the group consisting of polyvinyl pyrrolidone, starch and gelatin.

[0080] The crumbly mass is then sieved to select for a range of granule sizes. The size of the granules selected by the sieving process may be in the range of 250-1700  $\mu\text{m}$ , more preferably 1000-1700  $\mu\text{m}$ , most preferably 500-1000  $\mu\text{m}$ . The sieved granules are then dried at 90-110°C, more preferably at 105°C.

[0081] The dried granules are then heated to 700-800°C to remove the pore-forming agent. The temperature is then raised to 1000-1200°C, more preferably 1150°C, for sintering. The sintered granules undergo a slow cooling procedure to attain pure crystalline  $\beta$ -TCP. In a preferred embodiment the temperature is lowered from 1150°C to 39°C in 6 hours. After sintering, weight loss and shrinkage takes place in the sample. Pores are formed in the TCP and the pores are surrounded by the

- 29 -

skeleton of sintered TCP. The sintered granules are resieved using the same size sieve as previously used and mixed with a binder as previously described to form a moldable putty composition.

5 [0082] Alternatively, the porous  $\beta$ -TCP granules may be prepared by mixing the TCP powder with the pore-forming agent. The mixture is blended to achieve homogeneity and pressed into slugs using a press, rotary tablet machine or chilsonators. The slugs are heated to 700-800°C to  
10 remove the pore-forming agent and sintered at 1000-1100°C, preferably at 1150°C. The porous slugs are then fractured into the appropriate particle size range of 250-1700  $\mu\text{m}$ , more preferably 1000-1700  $\mu\text{m}$ , and most preferably 500-1000  $\mu\text{m}$ . The porous granules are then  
15 mixed with a binder to form a moldable putty composition.

#### Moldable Putty Composition

[0083] The porous  $\beta$ -TCP of this invention may be combined with a biocompatible binder to form a moldable putty composition. The moldable putty may be in the form  
20 of a paste or a semi-solid having sufficient viscosity. The moldable putty composition enables the positioning and shaping within the voids, defects or other areas in which new bone growth is desired. The cohesiveness of the putty also prevents the problems of particle  
25 migration associated with grafting materials for orthopedic, maxillofacial and dental applications.

[0084] The binder according to this invention must be biodegradable, biocompatible and have fluid flow properties. The binders contemplated as useful herein  
30 include, but are not limited to: art-recognized suspending agents, viscosity-producing agents, gel-forming agents and emulsifying agents. Other candidates are agents used to suspend ingredients for topical, oral

- 30 -

or parental administration. Yet other candidates are agents useful as tablet binders, disintegrants or emulsion stabilizers. Still other candidates are agents used in cosmetics, toiletries and food products.

- 5 Reference manuals such as the USP XXII -NF XVII (The Nineteen Ninety U.S. Pharmacopeia and the National Formulary (1990)) categorize and describe such agents. Preferred binders include resorbable macromolecules from biological or synthetic sources including sodium
- 10 alginate, hyaluronic acid, cellulose derivatives such as alkylcelluloses including methylcellulose, carboxy methylcellulose, carboxy methylcellulose sodium, carboxy methylcellulose calcium or other salts, hydroxy
- 15 alkylcelluloses including hydroxypropyl methylcellulose, hydroxybutyl methylcellulose, hydroxyethyl methylcellulose, hydroxyethyl cellulose, alkylhydroxyalkyl celluloses including methylhydroxyethyl cellulose, collagen, peptides, mucin, chondroitin sulfate and the like.
- 20 [0085] Carboxymethylcellulose (CMC) sodium is a preferred binder. CMC is commercially available from suppliers such as, but not limited to: Hercules Inc., Aqualon® Division, Delaware; FMC Corporation, Pennsylvania; British Celanese, Ltd., United Kingdom; and
- 25 Henkel KGaA, United Kingdom. Carboxymethylcellulose sodium is the sodium salt of a polycarboxymethyl ether of cellulose with a typical molecular weight ranging from 90,000 -700,000. Various grades of
- 30 carboxymethylcellulose sodium are commercially available which have differing viscosities. Viscosities of various grades of carboxymethylcellulose sodium are reported in Handbook of Pharmaceutical Excipients (2nd Edition), American Pharmaceutical Association & Royal Pharmaceutical Society of Great Britain. For example,

- 31 -

low viscosity 50-200 cP, medium viscosity 400-800 cP, high viscosity 1500-3000 cP. A number of grades of carboxymethylcellulose sodium are commercially available, the most frequently used grade having a degree of substitution (DS) of 0.7. The DS is defined as the average number of hydroxyl groups substituted per anhydroglucose unit. It is this DS which determines the aqueous solubility of the polymer. The degree of substitution and the standard viscosity of an aqueous solution of stated concentration is indicated on any carboxymethylcellulose sodium labeling. Low viscosity CMC (Aqualon® Division, Hercules, Inc., Wilmington, DE) is currently preferred. The currently preferred degrees of substitution range from 0.65-0.90 (DS=0.7, Aqualon® Type 7L).

[0086] Aside from binders that are flowable at room temperature, binders also include reagents such as gelatin, that are solubilized in warm or hot aqueous solutions, and are transformed into a non-flowable gel upon cooling. The gelatin composition is formulated so that the composition is flowable at temperatures above the body temperature of the mammal for implant, but transitions to relatively non-flowable gel at or slightly above such body temperature.

[0087] In one embodiment, the binder of this invention is selected from a class of high molecular weight hydrogels including sodium hyaluronate (~ 500-3000 kD), chitosan (~ 100 - 300 kD), poloxamer (~ 7 - 18 kD), and glycosaminoglycan (~ 2000 - 3000 kD). In a preferred embodiment, the glycosaminoglycan is N,O-carboxymethylchitosan glucosamine. Hydrogels are cross-linked hydrophilic polymers in the form of a gel which have a three-dimensional network. Hydrogel matrices can carry a net positive or net negative charge,

- 32 -

or may be neutral. A typical net negative charged matrix is alginate. Hydrogels carrying a net positive charge may be typified by extracellular matrix components such as collagen and laminin. Examples of commercially available extracellular matrix components include Matrigel™ (Dulbecco's modified eagle's medium with 50 µg/ml gentamicin) and Vitrogen™ (a sterile solution of purified, pepsin-solubilized bovine dermal collagen dissolved in 0.012 N HCL). An example of a net neutral hydrogel is highly crosslinked polyethylene oxide, or polyvinylalcohol.

[0088] In another embodiment, the binder of this invention may also be selected from a class of polymers selected from the group comprising polylactic acid, polyglycolic acid, co-polymers of polylactic acid and polyglycolic acid, polyhydroxybutyric acid, polymalic acid, polyglutamic acid, and polylactone. In order to have gradual polymer replacement in the material by *in situ* tissue ingrowth over a several-day to several-week period, the molecular weight of the polymer should be compatible with the required degradation rate of the polymer.

[0089] In another preferred embodiment, the binder is polyethylene glycol. A mixture of low- and high-molecular-weight polyethylene glycols can produce a paste with the proper viscosity. For example, a mixture of polyethylene glycols of molecular weight 400-600 daltons and 1500 daltons at the proper ratio would be effective.

[0090] In yet another embodiment, the binder is selected from a class of polysaccharides with an average molecular weight of about 200,000 to 5,000,000 daltons consisting of dextran, dextran sulfate, diethylaminoethyl dextran, dextran phosphate or mixtures thereof. Lower molecular weight polysaccharides have the advantage of a

- 33 -

faster dextran absorption rate, resulting in earlier exposure of the porous  $\beta$ -TCP material. If it is desired that dextrans remain in the site for an extended period, dextrans of relatively high molecular weight may be used.

5 Other preferred polysaccharides include starch, fractionated starch, amylopectin, agar, gum arabic, pullullan, agarose, carrageenan, dextrans, fructans, inulin, mannans, xylans, arabinans, glycogens, glucans, xanthan gum, guar gum, locust bean gum, tragacanth gum,  
10 karaya gum, and derivatives and mixtures thereof.

[0091] In another preferred embodiment, the binder is selected from the group consisting of mannitol, white petrolatum, mannitol/dextran combinations, mannitol/white petrolatum combinations, sesame oil, fibrin glue and  
15 admixtures thereof. Fibrin glue is currently a preferred binder, which comprises a mixture of mammalian fibrinogen and thrombin. Human fibrinogen is commercially available in products such as, but not limited to Tissucol<sup>®</sup> (Immuno AG, Vienna, Austria), Beriplast<sup>®</sup> (Behringwerke, Marburg,  
20 Germany), Biocoll<sup>®</sup> (Centre de Transfusion Sanguine de Lille, Pours, France) and Transglutine<sup>®</sup> (CNTS Fractionation Centre, Strasbourg, France). Fibrin glue may also be made of fibrinogen and thrombin from other mammalian sources, such as, for example, bovine and  
25 murine sources.

[0092] It is preferred that the binder is selected from the group consisting of sodium alginate, hyaluronic acid, sodium hyaluronate, gelatin, collagen, peptides, mucin, chondroitin sulfate, chitosan, poloxamer,  
30 glycosaminoglycan, polysaccharide, polyethylene glycol, methylcellulose, carboxy methylcellulose, carboxy methylcellulose sodium, carboxy methylcellulose calcium, hydroxypropyl methylcellulose, hydroxybutyl methylcellulose, hydroxyethyl methylcellulose,

- 34 -

hydroxyethylcellulose, methylhydroxyethyl cellulose, hydroxyethyl cellulose, polylactic acid, polyglycolic acid, co-polymers of polylactic acid and polyglycolic acid, polyhydroxybutyric acid, polymalic acid, polyglutamic acid, polylactone, mannitol, white petrolatum, mannitol/dextran combinations, mannitol/white petrolatum combinations, sesame oil, fibrin glue and admixtures thereof.

[0093] More preferably, the binder is selected from the group consisting of sodium alginate, hyaluronic acid, methylcellulose, carboxy methylcellulose, carboxy methylcellulose sodium, carboxy methylcellulose calcium, hydroxypropyl methylcellulose, hydroxybutyl methylcellulose, hydroxyethyl methylcellulose, hydroxyethylcellulose, methylhydroxyethyl cellulose, hydroxyethyl cellulose and admixtures thereof. Most preferably, the binder is selected from the group consisting of sodium alginate, hyaluronic acid, carboxy methylcellulose, carboxy methylcellulose sodium and carboxy methylcellulose calcium.

[0094] The minimum amount of binder is the amount necessary to give easy formability and provide sufficient particle cohesion and shape retention during the period of tissue ingrowth. In one embodiment, the weight ratio of porous  $\beta$ -TCP to carboxy methylcellulose sodium is in the range of 1:0.1 to 1:1.25. In a preferred embodiment, the ratio of porous  $\beta$ -TCP to CMC sodium is 1:0.4.

[0095] The invention also relates to a kit for bone implant comprising the porous  $\beta$ -TCP material of the invention and at least one additional bioactive agent selected from the group consisting of bone morphogenic proteins and antibiotics. The kit comprising the porous  $\beta$ -TCP material and a bone morphogenic protein may further comprise a morphogenic protein stimulatory factor. In



- 35 -

one embodiment, the kit further comprises a binder. In another embodiment, the kit comprises the porous  $\beta$ -TCP material of the invention and a binder.

#### Bone Morphogenic Protein Family

5 [0096] The BMP family, named for its representative bone morphogenic/osteogenic protein family members, belongs to the TGF- $\beta$  protein superfamily. Of the reported "BMPs" (BMP-1 to BMP-18), isolated primarily based on sequence homology, all but BMP-1 remain  
10 classified as members of the BMP family of morphogenic proteins (Ozkaynak et al., EMBO J., 9, pp. 2085-93 (1990)).

[0097] The BMP family includes other structurally-related members which are morphogenic proteins, including  
15 the *drosophila* decapentaplegic gene complex (DPP) products, the Vg1 product of *Xenopus laevis* and its murine homolog, Vgr-1 (see, e.g., Massagué, Annu. Rev. Cell Biol., 6, pp. 597-641 (1990), incorporated herein by reference).

20 [0098] The C-terminal domains of BMP-3, BMP-5, BMP-6, and OP-1 (BMP-7) are about 60% identical to that of BMP-2, and the C-terminal domains of BMP-6 and OP-1 are 87% identical. BMP-6 is likely the human homolog of the murine Vgr-1 (Lyons et al., Proc. Natl. Acad. Sci. U.S.A., 86, pp. 4554-59 (1989)); the two proteins are 92%  
25 identical overall at the amino acid sequence level (U. S. Patent No. 5,459,047, incorporated herein by reference). BMP-6 is 58% identical to the *Xenopus* Vg-1 product.

#### Biochemical Structural and Functional Properties of Bone Morphogenic Proteins

30 [0099] The naturally occurring bone morphogens share substantial amino acid sequence homology in their C-

- 36 -

terminal regions (domains). Typically, the above-mentioned naturally occurring osteogenic proteins are translated as a precursor, having an N-terminal signal peptide sequence typically less than about 30 residues, followed by a "pro" domain that is cleaved to yield the mature C-terminal domain of approximately 100-140 amino acids. The signal peptide is cleaved rapidly upon translation, at a cleavage site that can be predicted in a given sequence using the method of Von Heijne Nucleic Acids Research, 14, pp. 4683-4691 (1986). The pro domain typically is about three times larger than the fully processed mature C-terminal domain.

[0100] Another characteristic of the BMP protein family members is their apparent ability to dimerize. Several bone-derived osteogenic proteins (OPs) and BMPs are found as homo- and heterodimers in their active forms. The ability of OPs and BMPs to form heterodimers may confer additional or altered morphogenic inductive capabilities on morphogenic proteins. Heterodimers may exhibit qualitatively or quantitatively different binding affinities than homodimers for OP and BMP receptor molecules. Altered binding affinities may in turn lead to differential activation of receptors that mediate different signaling pathways, which may ultimately lead to different biological activities or outcomes. Altered binding affinities could also be manifested in a tissue or cell type-specific manner, thereby inducing only particular progenitor cell types to undergo proliferation and/or differentiation.

[0101] In preferred embodiments, the pair of morphogenic polypeptides have amino acid sequences each comprising a sequence that shares a defined relationship with an amino acid sequence of a reference morphogen. Herein, preferred osteogenic polypeptides share a defined

- 37 -

relationship with a sequence present in osteogenically active human OP-1, SEQ ID NO: 1. However, any one or more of the naturally occurring or biosynthetic sequences disclosed herein similarly could be used as a reference sequence. Preferred osteogenic polypeptides share a defined relationship with at least the C-terminal six cysteine domain of human OP-1, residues 335-431 of SEQ ID NO: 1. Preferably, osteogenic polypeptides share a defined relationship with at least the C-terminal seven cysteine domain of human OP-1, residues 330-431 of SEQ ID NO: 1. That is, preferred polypeptides in a dimeric protein with bone morphogenic activity each comprise a sequence that corresponds to a reference sequence or is functionally equivalent thereto.

[0102] Functionally equivalent sequences include functionally equivalent arrangements of cysteine residues disposed within the reference sequence; including amino acid insertions or deletions which alter the linear arrangement of these cysteines, but do not materially impair their relationship in the folded structure of the dimeric morphogen protein, including their ability to form such intra- or inter-chain disulfide bonds as may be necessary for morphogenic activity. Functionally equivalent sequences further include those wherein one or more amino acid residues differs from the corresponding residue of a reference sequence, e.g., the C-terminal seven cysteine domain (also referred to herein as the conserved seven cysteine skeleton) of human OP-1, provided that this difference does not destroy bone morphogenic activity. Accordingly, conservative substitutions of corresponding amino acids in the reference sequence are preferred. Amino acid residues that are conservative substitutions for corresponding residues in a reference sequence are those that are

- 38 -

physically or functionally similar to the corresponding reference residues, e.g., that have similar size, shape, electric charge, chemical properties including the ability to form covalent or hydrogen bonds, or the like.

5. Particularly preferred conservative substitutions are those fulfilling the criteria defined for an accepted point mutation in Dayhoff et al., *supra*, the teachings of which are incorporated by reference herein.

[0103] Conservative substitutions typically include the substitution of one amino acid for another with similar characteristics, e.g., substitutions within the following groups: valine, glycine; glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. The term "conservative variation" also includes the use of a substituted amino acid in place of an unsubstituted parent amino acid provided that antibodies raised to the substituted polypeptide also immunoreact with the unsubstituted polypeptide.

[0104] The osteogenic protein OP-1 has been described (see, e.g., Oppermann et al., U. S. Patent No. 5,354,557, incorporated herein by reference). Natural-sourced osteogenic protein in its mature, native form is a glycosylated dimer typically having an apparent molecular weight of about 30-36 kDa as determined by SDS-PAGE. When reduced, the 30 kDa protein gives rise to two glycosylated peptide subunits having apparent molecular weights of about 16 kDa and 18 kDa. In the reduced state, the protein has no detectable osteogenic activity. The unglycosylated protein, which also has osteogenic activity, has an apparent molecular weight of about 27 kDa. When reduced, the 27 kDa protein gives rise to two unglycosylated polypeptides, having molecular weights of

- 39 -

about 14 kDa to 16 kDa, capable of inducing endochondral bone formation in a mammal. Osteogenic proteins may include forms having varying glycosylation patterns, varying N-termini, and active truncated or mutated forms of native protein. As described above, particularly useful sequences include those comprising the C-terminal 96 or 102 amino acid sequences of DPP (from *Drosophila*), Vg1 (from *Xenopus*), Vgr-1 (from mouse), the OP-1 and OP-2 proteins, (see U.S. Pat. No. 5,011,691 and Oppermann et al., incorporated herein by reference), as well as the proteins referred to as BMP-2, BMP-3, BMP-4 (see WO88/00205, U.S. Patent No. 5,013,649 and WO91/18098, incorporated herein by reference), BMP-5 and BMP-6 (see WO90/11366, PCT/US90/01630, incorporated herein by reference), BMP-8 and BMP-9.

[0105] Preferred morphogenic and osteogenic proteins of this invention comprise at least one polypeptide selected from the group consisting of OP-1 (BMP-7), OP-2, OP-3, COP-1, COP-3, COP-4, COP-5, COP-7, COP-16, BMP-2, BMP-3, BMP-3b, BMP-4, BMP-5, BMP-6, BMP-9, BMP-10, BMP-11, BMP-12, BMP-13, BMP-14, BMP-15, BMP-16, BMP-17, BMP-18, GDF-1, GDF-2, GDF-3, GDF-5, GDF-6, GDF-7, GDF-8, GDF-9, GDF-10, GDF-11, GDF-12, MP121, dorsalin-1, DPP, Vg-1, Vgr-1, 60A protein, NODAL, UNIVIN, SCREW, ADMP, NEURAL, TGF- $\beta$  and amino acid sequence variants and homologs thereof, including species homologs, thereof. Preferably, the morphogenic protein comprises at least one polypeptide selected from the group consisting of OP-1 (BMP-7), BMP-2, BMP-4, BMP-5 and BMP-6; more preferably, OP-1 (BMP-7) and BMP-2; and most preferably, OP-1 (BMP-7).

[0106] Publications disclosing these sequences, as well as their chemical and physical properties, include: OP-1 and OP-2 (U.S. Patent No. 5,011,691; U.S. Patent

- 40 -

No. 5,266,683; Ozkaynak et al., EMBO J., 9, pp. 2085-2093 (1990); OP-3 (WO94/10203 (PCT US93/10520)), BMP-2, BMP-3, BMP-4, (WO88/00205; Wozney et al. Science, 242, pp. 1528-1534 (1988)), BMP-5 and BMP-6, (Celeste et al., PNAS, 87, 9843-9847 (1991)), Vgr-1 (Lyons et al., PNAS, 86, pp. 4554-4558 (1989)); DPP (Padgett et al. Nature, 325, pp. 81-84 (1987)); Vg-1 (Weeks, Cell, 51, pp. 861-867 (1987)); BMP-9 (WO95/33830 (PCT/US95/07084)); BMP-10 (WO94/26893 (PCT/US94/05290)); BMP-11 (WO94/26892 (PCT/US94/05288)); BMP-12 (WO95/16035 (PCT/US94/14030)); BMP-13 (WO95/16035 (PCT/US94/14030)); GDF-1 (WO92/00382 (PCT/US91/04096) and Lee et al. PNAS, 88, pp. 4250-4254 (1991); GDF-8 (WO94/21681 (PCT/US94/03019)); GDF-9 (WO94/15966 (PCT/US94/00685)); GDF-10 (WO95/10539 (PCT/US94/11440)); GDF-11 (WO96/01845 (PCT/US95/08543)); BMP-15 (WO96/36710 (PCT/US96/06540)); MP-121 (WO96/01316 (PCT/EP95/02552); GDF-5 (CDMP-1, MP52) (WO94/15949 (PCT/US94/00657) and WO96/14335 (PCT/US94/12814) and WO93/16099 (PCT/EP93/00350)); GDF-6 (CDMP-2, BMP13) (WO95/01801 (PCT/US94/07762) and WO96/14335 and WO95/10635 (PCT/US94/14030)); GDF-7 (CDMP-3, BMP12) (WO95/10802 (PCT/US94/07799) and WO95/10635 (PCT/US94/14030)) The above publications are incorporated herein by reference. In another embodiment, useful proteins include biologically active biosynthetic constructs, including novel biosynthetic morphogenic proteins and chimeric proteins designed using sequences from two or more known morphogens.

[0107] In another embodiment of this invention, a morphogenic protein may be prepared synthetically for use in concert with a MPSF to induce tissue formation. Morphogenic proteins prepared synthetically may be native, or may be non-native proteins, i.e., those not otherwise found in nature.

- 41 -

[0108] Non-native osteogenic proteins have been synthesized using a series of consensus DNA sequences (U.S. Patent No. 5,324,819, incorporated herein by reference). These consensus sequences were designed  
5 based on partial amino acid sequence data obtained from natural osteogenic products and on their observed homologies with other genes reported in the literature having a presumed or demonstrated developmental function.

[0109] Several of the biosynthetic consensus sequences  
10 (called consensus osteogenic proteins or "COPs") have been expressed as fusion proteins in prokaryotes. Purified fusion proteins may be cleaved, refolded, combined with at least one MPSF (optionally in a matrix or device), implanted in an established animal model and  
15 shown to have bone- and/or cartilage-inducing activity. The currently preferred synthetic osteogenic proteins comprise two synthetic amino acid sequences designated COP-5 (SEQ. ID NO: 2) and COP-7 (SEQ. ID NO: 3)

[0110] Oppermann et al., U. S. Patent Nos. 5,011,691  
20 and 5,324,819, which are incorporated herein by reference, describe the amino acid sequences of COP-5 and COP-7 as shown below:

	COP5	LYVDFS-DVGWDDWIVAPPGYQAFYCHGECPFPLAD
	COP7	LYVDFS-DVGWNDWIVAPPGYHAFYCHGECPFPLAD
25	COP5	HFNSTN--H-AVVQTLVNSVNSKI--PKACCVPTELSA
	COP7	HLNSTN--H-AVVQTLVNSVNSKI--PKACCVPTELSA
	COP5	ISMLYLDENEKVVLKYNQEMVVEGCGCR
	COP7	ISMLYLDENEKVVLKYNQEMVVEGCGCR

[0111] In these amino acid sequences, the dashes (-)  
30 are used as fillers only to line up comparable sequences

- 42 -

in related proteins. Differences between the aligned amino acid sequences are highlighted.

[0112] The DNA and amino acid sequences of these and other BMP family members are published and may be used by those of skill in the art to determine whether a newly identified protein belongs to the BMP family. New BMP-related gene products are expected by analogy to possess at least one morphogenic activity and thus classified as a BMP.

[0113] In one preferred embodiment of this invention, the morphogenic protein comprises a pair of subunits disulfide bonded to produce a dimeric species, wherein at least one of the subunits comprises a recombinant peptide belonging to the BMP protein family. In another preferred embodiment of this invention, the morphogenic protein comprises a pair of subunits that produce a dimeric species formed through non-covalent interactions, wherein at least one of the subunits comprises a recombinant peptide belonging to the BMP protein family. Non-covalent interactions include Van der Waals, hydrogen bond, hydrophobic and electrostatic interactions. The dimeric species may be a homodimer or heterodimer and is capable of inducing cell proliferation and/or tissue formation.

[0114] In certain preferred embodiments, bone morphogenic proteins useful herein include those in which the amino acid sequences comprise a sequence sharing at least 70% amino acid sequence homology or "similarity", and preferably 80% homology or similarity, with a reference morphogenic protein selected from the foregoing naturally occurring proteins. Preferably, the reference protein is human OP-1, and the reference sequence thereof is the C-terminal seven cysteine domain present in osteogenically active forms of human OP-1, residues 330-



- 43 -

431 of SEQ ID NO: 1. In certain embodiments, a polypeptide suspected of being functionally equivalent to a reference morphogen polypeptide is aligned therewith using the method of Needleman, et al., *supra*, implemented conveniently by computer programs such as the Align program (DNASTar, Inc.). As noted above, internal gaps and amino acid insertions in the candidate sequence are ignored for purposes of calculating the defined relationship, conventionally expressed as a level of amino acid sequence homology or identity, between the candidate and reference sequences. "Amino acid sequence homology" is understood herein to include both amino acid sequence identity and similarity. Homologous sequences share identical and/or similar amino acid residues, where similar residues are conservation substitutions for, or "allowed point mutations" of, corresponding amino acid residues in an aligned reference sequence. Thus, a candidate polypeptide sequence that shares 70% amino acid homology with a reference sequence is one in which any 70% of the aligned residues are either identical to, or are conservative substitutions of, the corresponding residues in a reference sequence. In a currently preferred embodiment, the reference sequence is OP-1. Bone morphogenic proteins useful herein accordingly include allelic, phylogenetic counterpart and other variants of the preferred reference sequence, whether naturally-occurring or biosynthetically produced (e.g., including "muteins" or "mutant proteins"), as well as novel members of the general morphogenic family of proteins, including those set forth and identified above. Certain particularly preferred morphogenic polypeptides share at least 60% amino acid identity with the preferred reference sequence of human OP-1, still more preferably at least 65% amino acid identity therewith.

- 44 -

[0115] In another embodiment, useful osteogenic proteins include those sharing the conserved seven cysteine domain and sharing at least 70% amino acid sequence homology (similarity) within the C-terminal active domain, as defined herein. In still another embodiment, the osteogenic proteins of the invention can be defined as osteogenically active proteins having any one of the generic sequences defined herein, including OPX (SEQ ID NO: 4) and Generic Sequences 7 (SEQ ID NO: 5) and 8 (SEQ ID NO: 6), or Generic Sequences 9 (SEQ ID NO: 7) and 10 (SEQ ID NO: 8).

[0116] The family of bone morphogenic polypeptides useful in the present invention, and members thereof, can be defined by a generic amino acid sequence. For example, Generic Sequence 7 (SEQ ID NO: 5) and Generic Sequence 8 (SEQ ID NO: 6) are 97 and 102 amino acid sequences, respectively, and accommodate the homologies shared among preferred protein family members identified to date, including at least OP-1, OP-2, OP-3, CBMP-2A, CBMP-2B, BMP-3, 60A, DPP, Vgl, BMP-5, BMP-6, Vgr-1, and GDF-1. The amino acid sequences for these proteins are described herein and/or in the art, as summarized above. The generic sequences include both the amino acid identity shared by these sequences in the C-terminal domain, defined by the six and seven cysteine skeletons (Generic Sequences 7 and 8, respectively), as well as alternative residues for the variable positions within the sequence. The generic sequences provide an appropriate cysteine skeleton where inter- or intramolecular disulfide bonds can form, and contain certain critical amino acids likely to influence the tertiary structure of the folded proteins. In addition, the generic sequences allow for an additional cysteine at position 36 (Generic Sequence 7) or position 41 (Generic

- 45 -

Sequence 8), thereby encompassing the morphogenically active sequences of OP-2 and OP-3.

Generic Sequence 7

				Leu	Xaa	Xaa	Xaa	Phe	Xaa	Xaa
				1				5		
5	Xaa	Gly	Trp	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Pro
			10					15		
	Xaa	Xaa	Xaa	Xaa	Ala	Xaa	Tyr	Cys	Xaa	Gly
			20					25		
	Xaa	Cys	Xaa	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Xaa
10			30					35		
	Xaa	Xaa	Xaa	Asn	His	Ala	Xaa	Xaa	Xaa	Xaa
			40					45		
	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
			50					55		
15	Xaa	Xaa	Xaa	Cys	Cys	Xaa	Pro	Xaa	Xaa	Xaa
			60					65		
	Xaa	Xaa	Xaa	Xaa	Xaa	Leu	Xaa	Xaa	Xaa	Xaa
			70					75		
	Xaa	Xaa	Xaa	Val	Xaa	Leu	Xaa	Xaa	Xaa	Xaa
20			80					85		
	Xaa	Met	Xaa	Val	Xaa	Xaa	Cys	Xaa	Cys	Xaa
			90					95		

wherein each Xaa independently is selected from a group of one or more specified amino acids defined as follows:

25 "res." means "residue" and Xaa at res.2 = (Tyr or Lys); Xaa at res.3 = Val or Ile); Xaa at res.4 = (Ser, Asp or Glu); Xaa at res.6 = (Arg, Gln, Ser, Lys or Ala); Xaa at res.7 = (Asp or Glu); Xaa at res.8 = (Leu, Val or Ile); Xaa at res. 11 = (Gln, Leu, Asp, His, Asn or Ser);

30 Xaa at res.12 = (Asp, Arg, Asn or Glu); Xaa at res.13 = (Trp or Ser); Xaa at res.14 = (Ile or Val); Xaa at res.15 = (Ile or Val); Xaa at res.16 (Ala or Ser); Xaa at res.18 = (Glu, Gln, Leu, Lys, Pro or Arg); Xaa at res.19 = (Gly or Ser); Xaa at res.20 = (Tyr or Phe); Xaa at res.21 =

35 (Ala, Ser, Asp, Met, His, Gln, Leu or Gly); Xaa at res.23 = (Tyr, Asn or Phe); Xaa at res.26 = (Glu, His, Tyr, Asp,

- 46 -

Gln, Ala or Ser); Xaa at res.28 = (Glu, Lys, Asp, Gln or Ala); Xaa at res.30 = (Ala, Ser, Pro, Gln, Ile or Asn); Xaa at res.31 = (Phe, Leu or Tyr); Xaa at res.33 = (Leu, Val or Met); Xaa at res.34 = (Asn, Asp, Ala, Thr or Pro);

5 Xaa at res.35 = (Ser, Asp, Glu, Leu, Ala or Lys); Xaa at res.36 = (Tyr, Cys, His, Ser or Ile); Xaa at res.37 = (Met, Phe, Gly or Leu); Xaa at res.38 = (Asn, Ser or Lys); Xaa at res.39 = (Ala, Ser, Gly or Pro); Xaa at res.40 = (Thr, Leu or Ser); Xaa at res.44 = (Ile, Val or Thr);

10 Xaa at res.45 = (Val, Leu, Met or Ile); Xaa at res.46 = (Gln or Arg); Xaa at res.47 = (Thr, Ala or Ser); Xaa at res.48 = (Leu or Ile); Xaa at res.49 = (Val or Met); Xaa at res.50 = (His, Asn or Arg); Xaa at res.51 = (Phe, Leu, Asn, Ser, Ala or Val); Xaa at res.52 =

15 (Ile, Met, Asn, Ala, Val, Gly or Leu); Xaa at res.53 = (Asn, Lys, Ala, Glu, Gly or Phe); Xaa at res.54 = (Pro, Ser or Val); Xaa at res.55 = (Glu, Asp, Asn, Gly, Val, Pro or Lys); Xaa at res.56 = (Thr, Ala, Val, Lys, Asp, Tyr, Ser, Gly, Ile or His); Xaa at res.57 = (Val, Ala or Ile);

20 Xaa at res.58 = (Pro or Asp); Xaa at res.59 = (Lys, Leu or Glu); Xaa at res.60 = (Pro, Val or Ala); Xaa at res.63 = (Ala or Val); Xaa at res.65 = (Thr, Ala or Glu); Xaa at res.66 = (Gln, Lys, Arg or Glu); Xaa at res.67 = (Leu, Met or Val); Xaa at res.68 = (Asn, Ser, Asp or Gly);

25 Xaa at res.69 = (Ala, Pro or Ser); Xaa at res.70 = (Ile, Thr, Val or Leu); Xaa at res.71 = (Ser, Ala or Pro); Xaa at res.72 = (Val, Leu, Met or Ile); Xaa at res.74 = (Tyr or Phe); Xaa at res.75 = (Phe, Tyr, Leu or His); Xaa at res.76 = (Asp, Asn or Leu); Xaa at res.77 =

30 (Asp, Glu, Asn, Arg or Ser); Xaa at res.78 = (Ser, Gln, Asn, Tyr or Asp); Xaa at res.79 = (Ser, Asn, Asp, Glu or Lys); Xaa at res.80 = (Asn, Thr or Lys); Xaa at res.82 = (Ile, Val or Asn); Xaa at res.84 = (Lys or Arg); Xaa at res.85 = (Lys, Asn, Gln, His, Arg or Val); Xaa at res.86

- 47 -

= (Tyr, Glu or His); Xaa at res.87 = (Arg, Gln, Glu or Pro); Xaa at res.88 = (Asn, Glu, Trp or Asp); Xaa at res.90 = (Val, Thr, Ala or Ile); Xaa at res.92 = (Arg, Lys, Val, Asp, Gln or Glu); Xaa at res.93 = (Ala, Gly, Glu or Ser); Xaa at res.95 = (Gly or Ala) and Xaa at res.97 = (His or Arg).

[0117] Generic Sequence 8 (SEQ ID NO: 6) includes all of Generic Sequence 7 and in addition includes the following sequence (SEQ ID NO: 9) at its N-terminus:

10		<u>SEQ ID NO: 9</u>			
	Cys	Xaa	Xaa	Xaa	Xaa
	1				5

[0118] Accordingly, beginning with residue 7, each "Xaa" in Generic Sequence 8 is a specified amino acid defined as for Generic Sequence 7, with the distinction that each residue number described for Generic Sequence 7 is shifted by five in Generic Sequence 8. Thus, "Xaa at res.2 = (Tyr or Lys)" in Generic Sequence 7 refers to Xaa at res. 7 in Generic Sequence 8. In Generic Sequence 8, Xaa at res.2 = (Lys, Arg, Ala or Gln); Xaa at res.3 = (Lys, Arg or Met); Xaa at res.4 = (His, Arg or Gln); and Xaa at res. 5 = (Glu, Ser, His, Gly, Arg, Pro, Thr, or Tyr).

[0119] In another embodiment, useful osteogenic proteins include those defined by Generic Sequences 9 and 10, defined as follows.

[0120] Specifically, Generic Sequences 9 and 10 are composite amino acid sequences of the following proteins: human OP-1, human OP-2, human OP-3, human BMP-2, human BMP-3, human BMP-4, human BMP-5, human BMP-6, human BMP-8, human BMP-9, human BMP 10, human BMP-11, Drosophila 60A, Xenopus Vg-1, sea urchin UNIVIN, human CDMP-1 (mouse

- 48 -

GDF-5), human CDMP-2 (mouse GDF-6, human BMP-13), human CDMP-3 (mouse GDF-7, human BMP-12), mouse GDF-3, human GDF-1, mouse GDF-1, chicken DORSALIN, dpp, Drosophila SCREW, mouse NODAL, mouse GDF-8, human GDF-8, mouse GDF-9, mouse GDF-10, human GDF-11, mouse GDF-11, human BMP-15, and rat BMP3b. Like Generic Sequence 7, Generic Sequence 9 is a 97 amino acid sequence that accommodates the C-terminal six cysteine skeleton and, like Generic Sequence 8, Generic Sequence 10 is a 102 amino acid sequence which accommodates the seven cysteine skeleton.

Generic Sequence 9

	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	
	1				5					10
	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Pro	Xaa	Xaa	Xaa
					15					20
15	Xaa	Xaa	Xaa	Xaa	Cys	Xaa	Gly	Xaa	Cys	Xaa
					25					30
	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
					35					40
	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
					45					50
	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
					55					60
	Xaa	Cys	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
					65					70
20	Xaa	Xaa	Leu	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
					75					80
	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
					85					90
	Xaa	Xaa	Xaa	Cys	Xaa	Cys	Xaa			
					95					

wherein each Xaa is independently selected from a group of one or more specified amino acids defined as follows:

"res." means "residue" and Xaa at res. 1 = (Phe, Leu or Glu); Xaa at res. 2 = (Tyr, Phe, His, Arg, Thr, Lys, Gln, Val or Glu); Xaa at res. 3 = (Val, Ile, Leu or Asp); Xaa

- 49 -

at res. 4 = (Ser, Asp, Glu, Asn or Phe); Xaa at res. 5 =  
(Phe or Glu); Xaa at res. 6 = (Arg, Gln, Lys, Ser, Glu,  
Ala or Asn); Xaa at res. 7 = (Asp, Glu, Leu, Ala or Gln);  
Xaa at res. 8 = (Leu, Val, Met, Ile or Phe); Xaa at res.  
5 9 = (Gly, His or Lys); Xaa at res. 10 = (Trp or Met); Xaa  
at res. 11 = (Gln, Leu, His, Glu, Asn, Asp, Ser or Gly);  
Xaa at res. 12 = (Asp, Asn, Ser, Lys, Arg, Glu or His);  
Xaa at res. 13 = (Trp or Ser); Xaa at res. 14 = (Ile or  
Val); Xaa at res. 15 = (Ile or Val); Xaa at res. 16 =  
10 (Ala, Ser, Tyr or Trp); Xaa at res. 18 = (Glu, Lys, Gln,  
Met, Pro, Leu, Arg, His or Lys); Xaa at res. 19 = (Gly,  
Glu, Asp, Lys, Ser, Gln, Arg or Phe); Xaa at res. 20 =  
(Tyr or Phe); Xaa at res. 21 = (Ala, Ser, Gly, Met, Gln,  
His, Glu, Asp, Leu, Asn, Lys or Thr); Xaa at res. 22 =  
15 (Ala or Pro); Xaa at res. 23 = (Tyr, Phe, Asn, Ala or  
Arg); Xaa at res. 24 = (Tyr, His, Glu, Phe or Arg); Xaa  
at res. 26 = (Glu, Asp, Ala, Ser, Tyr, His, Lys, Arg, Gln  
or Gly); Xaa at res. 28 = (Glu, Asp, Leu, Val, Lys, Gly,  
Thr, Ala or Gln); Xaa at res. 30 = (Ala, Ser, Ile, Asn,  
20 Pro, Glu, Asp, Phe, Gln or Leu); Xaa at res. 31 = (Phe,  
Tyr, Leu, Asn, Gly or Arg); Xaa at res. 32 = (Pro, Ser,  
Ala or Val); Xaa at res. 33 = (Leu, Met, Glu, Phe or  
Val); Xaa at res. 34 = (Asn, Asp, Thr, Gly, Ala, Arg, Leu  
or Pro); Xaa at res. 35 = (Ser, Ala, Glu, Asp, Thr, Leu,  
25 Lys, Gln or His); Xaa at res. 36 = (Tyr, His, Cys, Ile,  
Arg, Asp, Asn, Lys, Ser, Glu or Gly); Xaa at res. 37 =  
(Met, Leu, Phe, Val, Gly or Tyr); Xaa at res. 38 = (Asn,  
Glu, Thr, Pro, Lys, His, Gly, Met, Val or Arg); Xaa at  
res. 39 = (Ala, Ser, Gly, Pro or Phe); Xaa at res. 40 =  
30 (Thr, Ser, Leu, Pro, His or Met); Xaa at res. 41 = (Asn,  
Lys, Val, Thr or Gln); Xaa at res. 42 = (His, Tyr or  
Lys); Xaa at res. 43 = (Ala, Thr, Leu or Tyr); Xaa at  
res. 44 = (Ile, Thr, Val, Phe, Tyr, Met or Pro); Xaa at  
res. 45 = (Val, Leu, Met, Ile or His); Xaa at res. 46 =

- 50 -

(Gln, Arg or Thr); Xaa at res. 47 = (Thr, Ser, Ala, Asn or His); Xaa at res. 48 = (Leu, Asn or Ile); Xaa at res. 49 = (Val, Met, Leu, Pro or Ile); Xaa at res. 50 = (His, Asn, Arg, Lys, Tyr or Gln); Xaa at res. 51 = (Phe, Leu, Ser, Asn, Met, Ala, Arg, Glu, Gly or Gln); Xaa at res. 52 = (Ile, Met, Leu, Val, Lys, Gln, Ala or Tyr); Xaa at res. 53 = (Asn, Phe, Lys, Glu, Asp, Ala, Gln, Gly, Leu or Val); Xaa at res. 54 = (Pro, Asn, Ser, Val or Asp); Xaa at res. 55 = (Glu, Asp, Asn, Lys, Arg, Ser, Gly, Thr, Gln, Pro or His); Xaa at res. 56 = (Thr, His, Tyr, Ala, Ile, Lys, Asp, Ser, Gly or Arg); Xaa at res. 57 = (Val, Ile, Thr, Ala, Leu or Ser); Xaa at res. 58 = (Pro, Gly, Ser, Asp or Ala); Xaa at res. 59 = (Lys, Leu, Pro, Ala, Ser, Glu, Arg or Gly); Xaa at res. 60 = (Pro, Ala, Val, Thr or Ser); Xaa at res. 61 = (Cys, Val or Ser); Xaa at res. 63 = (Ala, Val or Thr); Xaa at res. 65 = (Thr, Ala, Glu, Val, Gly, Asp or Tyr); Xaa at res. 66 = (Gln, Lys, Glu, Arg or Val); Xaa at res. 67 = (Leu, Met, Thr or Tyr); Xaa at res. 68 = (Asn, Ser, Gly, Thr, Asp, Glu, Lys or Val); Xaa at res. 69 = (Ala, Pro, Gly or Ser); Xaa at res. 70 = (Ile, Thr, Leu or Val); Xaa at res. 71 = (Ser, Pro, Ala, Thr, Asn or Gly); Xaa at res. 2 = (Val, Ile, Leu or Met); Xaa at res. 74 = (Tyr, Phe, Arg, Thr, Tyr or Met); Xaa at res. 75 = (Phe, Tyr, His, Leu, Ile, Lys, Gln or Val); Xaa at res. 76 = (Asp, Leu, Asn or Glu); Xaa at res. 77 = (Asp, Ser, Arg, Asn, Glu, Ala, Lys, Gly or Pro); Xaa at res. 78 = (Ser, Asn, Asp, Tyr, Ala, Gly, Gln, Met, Glu, Asn or Lys); Xaa at res. 79 = (Ser, Asn, Glu, Asp, Val, Lys, Gly, Gln or Arg); Xaa at res. 80 = (Asn, Lys, Thr, Pro, Val, Ile, Arg, Ser or Gln); Xaa at res. 81 = (Val, Ile, Thr or Ala); Xaa at res. 82 = (Ile, Asn, Val, Leu, Tyr, Asp or Ala); Xaa at res. 83 = (Leu, Tyr, Lys or Ile); Xaa at res. 84 = (Lys, Arg, Asn, Tyr, Phe, Thr, Glu or Gly); Xaa at res. 85 = (Lys, Arg, His,



- 51 -

Gln, Asn, Glu or Val); Xaa at res. 86 = (Tyr, His, Glu or Ile); Xaa at res. 87 = (Arg, Glu, Gln, Pro or Lys); Xaa at res. 88 = (Asn, Asp, Ala, Glu, Gly or Lys); Xaa at res. 89 = (Met or Ala); Xaa at res. 90 = (Val, Ile, Ala, Thr, Ser or Lys); Xaa at res 91 = (Val or Ala); Xaa at res. 92 = (Arg, Lys, Gln, Asp, Glu, Val, Ala, Ser or Thr); Xaa at res. 93 = (Ala, Ser, Glu, Gly, Arg or Thr); Xaa at res. 95 = (Gly, Ala or Thr); Xaa at res. 97 = (His, Arg, Gly, Leu or Ser). Further, after res. 53 in rBMP3b and mGDF-10 there is an Ile; after res. 54 in GDF-1 there is a T ; after res. 54 in BMP3 there is a V; after res. 78 in BMP-8 and Dorsalin there is a G; after res. 37 in hGDF-1 there is Pro, Gly, Gly, Pro.

[0121] Generic Sequence 10 (SEQ ID NO: 8) includes all of Generic Sequence 9 (SEQ ID NO: 7) and in addition includes the following sequence (SEQ ID NO: 9) at its N-terminus:

SEQ ID NO: 9

Cys	Xaa	Xaa	Xaa	Xaa
1				5

[0122] Accordingly, beginning with residue 6, each "Xaa" in Generic Sequence 10 is a specified amino acid defined as for Generic Sequence 9, with the distinction that each residue number described for Generic Sequence 9 is shifted by five in Generic Sequence 10. Thus, "Xaa at res. 1 = ( Tyr, Phe, His, Arg, Thr, Lys, Gln, Val or Glu)" in Generic Sequence 9 refers to Xaa at res. 6 in Generic Sequence 10. In Generic Sequence 10, Xaa at res. 2 = (Lys, Arg, Gln, Ser, His, Glu, Ala, or Cys); Xaa at res. 3 = (Lys, Arg, Met, Lys, Thr, Leu, Tyr, or Ala); Xaa at res. 4 = (His, Gln, Arg, Lys, Thr, Leu, Val, Pro, or Tyr); and Xaa at res. 5 = (Gln, Thr, His, Arg, Pro, Ser, Ala, Gln, Asn, Tyr, Lys, Asp, or Leu).

- 52 -

[0123] As noted above, certain currently preferred bone morphogenic polypeptide sequences useful in this invention have greater than 60% identity, preferably greater than 65% identity, with the amino acid sequence defining the preferred reference sequence of hOP-1. These particularly preferred sequences include allelic and phylogenetic counterpart variants of the OP-1 and OP-2 proteins, including the Drosophila 60A protein. Accordingly, in certain particularly preferred embodiments, useful morphogenic proteins include active proteins comprising pairs of polypeptide chains within the generic amino acid sequence herein referred to as "OPX" (SEQ ID NO: 4), which defines the seven cysteine skeleton and accommodates the homologies between several identified variants of OP-1 and OP-2. As described therein, each Xaa at a given position independently is selected from the residues occurring at the corresponding position in the C-terminal sequence of mouse or human OP-1 or OP-2.

```

20  Cys Xaa Xaa His Glu Leu Tyr Val Ser Phe Xaa Asp Leu Gly Trp Xaa Asp Trp
    1           5           10          15
    Xaa Ile Ala Pro Xaa Gly Tyr Xaa Ala Tyr Tyr Cys Glu Gly Glu Cys Xaa Phe Pro
      20           25           30          35
25  Leu Xaa Ser Xaa Met Asn Ala Thr Asn His Ala Ile Xaa Gln Xaa Leu Val His Xaa
    40           45           50          55
    Xaa Xaa Pro Xaa Xaa Val Pro Lys Xaa Cys Cys Ala Pro Thr Xaa Leu Xaa Ala
      60           65           70
    Xaa Ser Val Leu Tyr Xaa Asp Xaa Ser Xaa Asn Val Ile Leu Xaa Lys Xaa Arg
    75           80           85          90
30  Asn Met Val Val Xaa Ala Cys Gly Cys His
    95           100

```

wherein Xaa at res. 2 = (Lys or Arg); Xaa at res. 3 = (Lys or Arg); Xaa at res. 11 = (Arg or Gln); Xaa at res. 16 = (Gln or Leu); Xaa at res. 19 = (Ile or Val); Xaa at res. 23 = (Glu or Gln); Xaa at res. 26 = (Ala or Ser); Xaa at res. 35 = (Ala or Ser); Xaa at res. 39 = (Asn or

- 53 -

Asp); Xaa at res. 41 = (Tyr or Cys); Xaa at res. 50 =  
(Val or Leu); Xaa at res. 52 = (Ser or Thr); Xaa at res.  
56 = (Phe or Leu); Xaa at res. 57 = (Ile or Met); Xaa at  
res. 58 = (Asn or Lys); Xaa at res. 60 = (Glu, Asp  
5 or Asn); Xaa at res. 61 = (Thr, Ala or Val); Xaa at res.  
65 = (Pro or Ala); Xaa at res. 71 = (Gln or Lys); Xaa at  
res. 73 = (Asn or Ser); Xaa at res. 75 = (Ile or Thr);  
Xaa at res. 80 = (Phe or Tyr); Xaa at res. 82 = (Asp or  
Ser); Xaa at res. 84 = (Ser or Asn); Xaa at res. 89 =  
10 (Lys or Arg); Xaa at res. 91 = (Tyr or His); and Xaa at  
res. 97 = (Arg or Lys).

[0124] In still another preferred embodiment, useful  
osteogenically active proteins have polypeptide chains  
with amino acid sequences comprising a sequence encoded  
15 by a nucleic acid that hybridizes, under low, medium or  
high stringency hybridization conditions, to DNA or RNA  
encoding reference morphogen sequences, e.g., C-terminal  
sequences defining the conserved seven cysteine domains  
of OP-1, OP-2, BMP-2, BMP-4, BMP-5, BMP-6, 60A, GDF-3,  
20 GDF-6, GDF-7 and the like. As used herein, high  
stringent hybridization conditions are defined as  
hybridization according to known techniques in 40%  
formamide, 5 X SSPE, 5 X Denhardt's Solution, and 0.1%  
SDS at 37°C overnight, and washing in 0.1 X SSPE, 0.1%  
25 SDS at 50°C. Standard stringent conditions are well  
characterized in commercially available, standard  
molecular cloning texts. See, for example, Molecular  
Cloning A Laboratory Manual, 2nd Ed., ed. by Sambrook,  
Fritsch and Maniatis (Cold Spring Harbor Laboratory  
30 Press: 1989); DNA Cloning, Volumes I and II (D.N. Glover  
ed., 1985); Oligonucleotide Synthesis (M.J. Gait ed.,  
1984); Nucleic Acid Hybridization (B. D. Hames & S.J.  
Higgins eds. 1984); and B. Perbal, A Practical Guide To

- 54 -

Molecular Cloning (1984), the disclosures of which are incorporated herein by reference.

[0125] As noted above, proteins useful in the present invention generally are dimeric proteins comprising a folded pair of the above polypeptides. Such morphogenic proteins are inactive when reduced, but are active as oxidized homodimers and when oxidized in combination with others of this invention to produce heterodimers. Thus, members of a folded pair of morphogenic polypeptides in a morphogenically active protein can be selected independently from any of the specific polypeptides mentioned above.

[0126] The bone morphogenic proteins useful in the materials and methods of this invention include proteins comprising any of the polypeptide chains described above, whether isolated from naturally-occurring sources, or produced by recombinant DNA or other synthetic techniques, and includes allelic and phylogenetic counterpart variants of these proteins, as well as muteins thereof, and various truncated and fusion constructs. Deletion or addition mutants also are envisioned to be active, including those which may alter the conserved C-terminal six or seven cysteine domain, provided that the alteration does not functionally disrupt the relationship of these cysteines in the folded structure. Accordingly, such active forms are considered the equivalent of the specifically described constructs disclosed herein. The proteins may include forms having varying glycosylation patterns, varying N-termini, a family of related proteins having regions of amino acid sequence homology, and active truncated or mutated forms of native or biosynthetic proteins, produced by expression of recombinant DNA in host cells.

- 55 -

[0127] The bone morphogenic proteins contemplated herein can be expressed from intact or truncated cDNA or from synthetic DNAs in prokaryotic or eukaryotic host cells, and purified, cleaved, refolded, and dimerized to form morphogenically active compositions. Currently preferred host cells include, without limitation, prokaryotes including *E. coli* or eukaryotes including yeast, or mammalian cells, such as CHO, COS or BSC cells. One of ordinary skill in the art will appreciate that other host cells can be used to advantage. Detailed descriptions of the bone morphogenic proteins useful in the practice of this invention, including how to make, use and test them for osteogenic activity, are disclosed in numerous publications, including U.S. Patent Nos. 5,266,683 and 5,011,691, the disclosures of which are incorporated by reference herein, as well as in any of the publications recited herein, the disclosures of which are incorporated herein by reference.

[0128] Thus, in view of this disclosure and the knowledge available in the art, skilled genetic engineers can isolate genes from cDNA or genomic libraries of various different biological species, which encode appropriate amino acid sequences, or construct DNAs from oligonucleotides, and then can express them in various types of host cells, including both prokaryotes and eukaryotes, to produce large quantities of active proteins capable of stimulating endochondral bone morphogenesis in a mammal.

#### Morphogenic Protein Stimulatory Factors (MPSF)

[0129] A morphogenic protein stimulatory factor (MPSF) according to this invention is a factor that is capable of stimulating the ability of a morphogenic protein to induce tissue formation from a progenitor cell. The MPSF

- 56 -

may have an additive effect on tissue induction by the morphogenic protein. Preferably, the MPSF has a synergistic effect on tissue induction by the morphogenic protein.

5     [0130]     The progenitor cell that is induced to proliferate and/or differentiate by the morphogenic protein of this invention is preferably a mammalian cell. Progenitor cells include mammalian chondroblasts, myoblasts, osteoblasts, neuroblasts and vascular tissue  
10   precursor cells, all earlier developmental precursors thereof, and all cells that develop therefrom (e.g., chondroblasts, pre-chondroblasts and chondrocytes). However, morphogenic proteins are highly conserved throughout evolution, and non-mammalian progenitor cells  
15   are also likely to be stimulated by same- or cross-species morphogenic proteins and MPSF combinations. It is thus envisioned that when schemes become available for implanting xenogeneic cells into humans without causing adverse immunological reactions, non-mammalian progenitor  
20   cells stimulated by morphogenic protein and a MPSF according to the procedures set forth herein will be useful for tissue regeneration and repair in humans.

   [0131]     One or more MPSFs are selected for use in concert with one or more morphogenic proteins according  
25   to the desired tissue type to be induced and the site at which the morphogenic protein and MPSF will be administered. The particular choice of a morphogenic protein(s)/MPSF(s) combination and the relative concentrations at which they are combined may be varied  
30   systematically to optimize the tissue type induced at a selected treatment site using the procedures described herein.

   [0132]     The preferred morphogenic protein stimulatory factors (MPSFs) of this invention are selected from the

- 57 -

group consisting of hormones, cytokines and growth factors. Most preferred MPSFs for inducing bone and/or cartilage formation in concert with an osteogenic protein comprise at least one compound selected from the group  
5 consisting of insulin-like growth factor I (IGF-I), estradiol, fibroblast growth factor (FGF), growth hormone (GH), growth and differentiation factor (GDF), hydrocortisone (HC), insulin, progesterone, parathyroid hormone (PTH), vitamin D (1,25-(OH)<sub>2</sub>D<sub>3</sub>), retinoic acid and  
10 an interleukin, particularly IL-6.

[0133] In another preferred embodiment of this invention, the MPSF comprises a compound or an agent that is capable of increasing the bioactivity of another MPSF. Agents that increase MPSF bioactivity include, for  
15 example, those that increase the synthesis, half-life, reactivity with other biomolecules such as binding proteins and receptors, or the bioavailability of the MPSF. These agents may comprise hormones, growth factors, peptides, cytokines, carrier molecules such as  
20 proteins or lipids, or other factors that increase the expression or the stability of the MPSF.

[0134] For example, when the selected MPSF is IGF-I, agents that increase its bioactivity include GH, PTH, vitamin D, and cAMP inducers, which may thus function as  
25 MPSFs according to this invention. In addition, almost all of the IGF-I in circulation and the extracellular space is bound by a group of high affinity binding proteins called IGFBPs which can augment or inhibit IGF-I  
30 bioactivity (see, e.g., Jones and Clemmons, Endocrine Reviews, 16, pp. 3-34 (1995)). Thus IGFBPs and agents which alter the levels of IGFBPs such that the bioactive IGF-I concentration is ultimately increased will also function as a MPSF according to this invention.

- 58 -

[0135] These or other agents that increase IGF-I bioactivity may be used alone as the primary MPSF, or one or more may be used as additional MPSFs in combination with IGF-I, to stimulate the tissue inductive activity of the morphogenic protein. One such preferred combination comprising at least two MPSFs for cartilage and bone formation is osteogenic protein OP-1, IGF-I and PTH.

[0136] Preferably, the MPSF is present in an amount capable of synergistically stimulating the tissue inductive activity of the morphogenic protein in a mammal. The relative concentrations of morphogenic protein and MPSF that will optimally induce tissue formation when administered to a mammal may be determined empirically by the skilled practitioner using the procedures described herein.

#### Implant Device

[0137] The invention also relates to an implant device for promoting bone formation, regeneration and repair. The implant device comprises the porous  $\beta$ -TCP material of the invention, and optionally at least one bioactive agent.

[0138] The implant device comprising the porous  $\beta$ -TCP material serves as a temporary scaffold and substratum for recruitment of migratory progenitor cells, and as a base for their subsequent anchoring and proliferation.

[0139] In a preferred embodiment, the implant device comprises the porous  $\beta$ -TCP matrix and a bioactive agent, which is dispersed or absorbed in the matrix. It is envisioned that the bioactive agent can include but is not limited to bone morphogenic proteins, growth factors such as EGF, PDGF, IGF, FGF, TGF- $\alpha$  and TGF- $\beta$ , cytokines, MPSF, hormones, peptides, lipids, trophic agents and therapeutic compositions including antibiotics and



- 59 -

chemotherapeutic agents, insulin, chemoattractant, chemotactic factors, enzymes, enzyme inhibitors. It is also envisioned that bioactive agents such as vitamins, cytoskeletal agents, autograft, allograft, cartilage fragments, living cells such as chondrocytes, bone marrow cells, mesenchymal stem cells, tissue transplants, immuno-suppressants may be added to the porous  $\beta$ -TCP.

[0140] The porous  $\beta$ -TCP matrix provides a sustained delivery or support system for the bioactive agent, which is released over time at the implantation site as the matrix material is slowly absorbed. In a preferred embodiment, the bioactive agent is encapsulated in the biodegradable agent. The resorption of the biodegradable agent and the gradual release of the bioactive agent provides a sustained release system. The dosage and rate of delivery of the bioactive agent may be controlled based on the nature of the porous matrix, the nature of the biodegradable agent and the nature of the binding interaction between the bioactive agent encapsulated in the biodegradable agent, the porous matrix and biodegradable agent. In a preferred embodiment, the bioactive agent is a bone morphogenic protein or a nucleic acid molecule that encodes BMP. In a most preferred embodiment, the BMP is OP-1.

[0141] In a preferred embodiment, the bioactive agent is a BMP. In a more preferred embodiment, the BMP is OP-1. The porous  $\beta$ -TCP matrix can protect the BMP and MPSF from non-specific proteolysis, and can accommodate each step of the cellular responses involved in progenitor cell induction during tissue development.

[0142] Studies have shown that the methodology for combining matrix and morphogenic proteins plays a role in achieving successful tissue induction. The optimal ratios of morphogenic protein to MPSF for a specific

- 60 -

combination and tissue type may be determined empirically by those of skill in the art. Greater amounts may be used for large implants. The procedures used to formulate BMP and MPSF into the matrix are sensitive to the physical and chemical state of both the proteins and the matrix.

[0143] In the preferred osteogenic device with porous  $\beta$ -TCP, the osteogenic protein diffuses out of the matrix into the implantation site and permits influx and efflux of cells. The osteogenic protein induces the progenitor cells to differentiate and proliferate. Progenitor cells may migrate into the matrix and differentiated cells can move out of the porous matrix into the implant site. The sequential cellular reactions in the interface of the bone matrix/osteogenic protein implants include: binding of fibrin and fibronectin to implanted matrix, migration and proliferation of mesenchymal cells, differentiation of the progenitor cells into chondroblasts, cartilage formation, cartilage calcification, vascular invasion, bone formation, remodeling, and bone marrow differentiation. The preferred osteogenic device with porous  $\beta$ -TCP material, can be applied to bone formation in various orthopedic, periodontal, and reconstructive procedures.

[0144] The implant device may also comprise a binder in an admixture with the bioactive agent and/or porous  $\beta$ -TCP material. The binder is added to form a moldable putty which may be shaped to fit a defect site or to take the form of a new tissue. The moldable putty composition can be held in place by the surrounding tissue or masticated muscle. It is preferred to shape the matrix to span a tissue defect and to take the desired form of the new tissue. In the case of bone repair of a non-union defect, for example, it is desirable to use

- 61 -

dimensions that span the non-union. Rat studies show that the new bone is formed essentially having the dimensions of the device implanted. Thus, the material may be used for subcutaneous or intramuscular implants.

5 In bone formation procedures, the material is slowly absorbed by the body and is replaced by bone in the shape of or very nearly the shape of the implant.

#### Prosthetic Device

[0145] It is also contemplated that the porous  $\beta$ -TCP material of the present invention may be used in a prosthetic device. The prosthetic device comprises a surface region that can be implanted adjacent to a target tissue of a mammal, and a composition that is disposed on the surface region. The prosthetic devices will be

15 useful for repairing orthopedic defects, injuries or anomalies in the treated mammal. Preferably, the mammal is a human patient. The prosthetic device may be made from a material comprising metal, ceramic or polymer composite material. Preferred devices comprise a load-

20 bearing core selected from Co-Cr-Mo alloys, titanium alloys and stainless steel. Preferred prosthetic devices are selected from the group consisting of a hip device, a fusion cage and a maxillofacial device.

[0146] The composition comprises the porous  $\beta$ -TCP material of the invention, and optionally, one or more agents selected from the group consisting of a bioactive agent or a binder dispersed in the porous  $\beta$ -TCP. In a preferred embodiment, the bioactive agent is encapsulated in the biodegradable agent. In a preferred embodiment,

25 the bioactive agent is a BMP or nucleic acid encoding BMP, more preferably, OP-1. Osteogenic protein-coated prosthetic devices may enhance the bond strength between the prosthesis and existing bone. (Rueger et al., U. S.

30

- 62 -

Patent No. 5,344,654, incorporated herein by reference). The composition may act as a coating for synthetically constructed bone material, such as for an artificial hip, replacement of diseased bone, correction of defects, or anchoring teeth. The composition is disposed on the surface of the implant in an amount sufficient to promote enhanced tissue growth into the surface. The amount of the composition sufficient to promote enhanced tissue growth may be determined empirically by those of skilled in the art using bioassays described in Rueger et al., U. S. Patent No. 5,344,654, incorporated herein by reference. Preferably, animal studies are performed to optimize the concentration of the composition components before a similar prosthetic device is used in the human patient.

[0147] In another preferred embodiment, the composition is applied to the clinical procedure of total joint arthroplasty in hips, knees, elbows and other joints, wherein a diseased or damaged natural joint is replaced by a prosthetic joint. For example, in a total hip arthroplasty, an acetabular cup is inserted with the composition in the acetabular socket of the pelvis to replace the natural acetabulum. The cup is held in place by the composition and secured by fixation screws.

Generally, the cavity or socket conforms to the outer surface of the acetabular cup. The composition can also be applied to total joint revision surgery, to strengthen the bondage between joint prosthetic devices and the bone.

[0148] In yet another preferred embodiment, the composition is applied to a clinical procedure called vertebroplasty. The composition is injected into the interior of a vertebral body. This method is used in the

- 63 -

treatment of osteoporosis to increase the density of bone.

[0149] In a preferred embodiment, the prosthetic device is selected from the group consisting of a fusion cage, a dowel and other devices having a pocket or chamber, such as an interbody fusion for containing the composition of the present invention. Preferably, the interbody fusion device is produced from material selected from the group consisting of titanium, PEEK (poly(etheretherketone)) and allograft. The interbody fusion in the cervical, thoracic and lumbar spine can be administered via an anterior or posterior approach. Alternatively, the composition of this invention can be used without an associated interbody device to achieve interbody fusion.

[0150] Spinal fusion cages are placed into the intervertebral space left after the removal of a damaged spinal disc to eliminate local motion and to participate in vertebral to vertebra bony fusion. As described in U.S. patent No. 5,015,247, incorporated herein by reference, the fusion cages are in the form of a cylindrical hollow member having an outside diameter larger than the space between two adjacent vertebrae to be fused. The interior space within the cylindrical hollow implant can be filled with the composition of this invention. The cylindrical implants can also include a threaded exterior to permit threaded insertion into a tapped bore formed in the adjacent vertebrae. Alternatively, some fusion implants have been designed to be impacted into the intradiscal space. As described in U.S. patent No. 6,146,420, incorporated herein by reference, the fusion device includes opposite end pieces with an integral central element. The central element has a much smaller diameter so that the fusion device

- 64 -

forms an annular pocket around the central element. The composition of this invention can be disposed within the annular pocket between the opposite end pieces.

5 [0151] In a preferred embodiment, the prosthetic device is used for repair of osseous and discoligamentous instability. The composition of this invention may be applied to the intervertebral area, resulting in superior fusion and consequently achieving definitive  
10 stabilization of a traumatized motor segment via a single dorsal approach. This application may eliminate the need to undergo a second operation for fractures of the thoracolumbar spine, which, at present, is often necessary but involves additional high risks. Also, this method avoids the problems associated with  
15 transplantation of autogenous cancellous bone and its associated risk of high morbidity might be avoided. See, e.g., Rueger et al., Orthopäde, 27, pp. 72-79 (1998).

[0152] In another preferred embodiment, the prosthetic device is a maxillofacial device. Maxillofacial devices  
20 are applied externally to correct facial defects resulting from cancer surgery, accidents, congenital deformities. In order to restore the masticatory deficiencies, a patient with marginal bone mass is first treated with the composition of this invention to pack  
25 and build up the surgical site. A maxillofacial anchoring and distracting system, as illustrated in U.S. patent No. 5,899,940, incorporated herein by reference, can be applied to increase the existing bone quality. Fixation devices, such as a standard threaded bone screw  
30 and simple pin point tack or self-locking and threaded bone tack screw device (U.S. patent No. 5,971,985, incorporated herein by reference), are used for the retention of tissue grafts and synthetic membranes to the maxillofacial bone graft site. Once the site has healed,

- 65 -

a second surgery is performed to insert the appropriate length endosseous dental implant and to restore masticatory function.

[0153] The invention also provides a method for promoting *in vivo* integration of an implantable prosthetic device of this invention into a target tissue of a mammal comprising the steps of a) providing on a surface of the prosthetic device a composition comprising the porous  $\beta$ -TCP material, optionally, at least one bioactive agent or a binder, and b) implanting the device in a mammal at a locus where the target tissue and the surface of the prosthetic device are maintained at least partially in contact for a time sufficient to permit tissue growth between the target tissue and the device.

15     Method of Inducing Bone Formation and Delivery

[0154] The invention also provides a method of inducing bone formation in a mammal. The mammal is preferably a human patient. The method comprises the step of implanting in the defect site of a mammal a composition comprising the porous  $\beta$ -TCP of the invention. In a preferred embodiment, the composition may further comprise a binder and/or a bioactive agent. The defect can be an endochondreal defect, an osteochondral defect or a segmental defect. The method can be applied to other defects which are not limited to, non-union fractures; bone cavities; tumor resection; fresh fractures (distracted or undistracted); cranial, maxillofacial and facial abnormalities, for example, in facial skeletal reconstruction, specifically, orbital floor reconstruction, augmentation of the alveolar ridge or sinus, periodontal defects and tooth extraction socket; cranioplasty, genioplasty, chin augmentation, palate reconstruction, and other large bony

- 66 -

reconstructions; vertebroplasty, interbody fusions in the cervical, thoracic and lumbar spine and posteriolateral fusions in the thoracic and lumbar spine; in osteomyelitis for bone regeneration; appendicular fusion, ankle fusion, total hip, knee and joint fusions or arthroplasty; correcting tendon and/or ligamentous tissue defects such as, for example, the anterior, posterior, lateral and medial ligaments of the knee, the patella and achilles tendons, and the like as well as those defects resulting from diseases such as cancer, arthritis, including osteoarthritis, and other bone degenerative disorders such as osteochondritis dessicans. The method may be used in bone augmentation, bone prosthesis, hard tissue implant, bone scaffolding, fixation systems (e.g. screws, sutures, suture anchors, staples, surgical tacks, clips, plates and screws).

[0155] The invention also provides a method of delivering a bioactive agent at a site requiring bone formation comprising the step of implanting the porous  $\beta$ -TCP and a bioactive agent at the defect site of a mammal. The method of delivering the bioactive agent may further include a binder. In a preferred embodiment, the bioactive agent is encapsulated in a biodegradable agent. In a preferred embodiment, the bioactive agent belongs to the bone morphogenic protein family. In another preferred embodiment, the bioactive agent is a nucleic acid molecule comprising a sequence encoding a BMP. Preferably, the nucleic acid is trapped in a carrier. In yet another embodiment, the bioactive agent is a bone cell or a cell transfected with nucleic acid encoding BMP. In another preferred embodiment, the delivery of the bioactive agent is sustained release. The biodegradable agent is preferably a biocompatible and non-immunogenic polymer, more preferably, PLGA. The bioactive agent is



- 67 -

preferably OP-1. The release rate of the bioactive agent can be controlled by altering the molecular weight of the PLGA. The degradation of PLGA commences when water penetrates the cement matrix to hydrolyze long polymer chains into short water soluble fragments. This results in a reduction in the molecular weight of the PLGA without loss in its physical properties. Gradually, further erosion of the polymer leads to the disruption of the polymer, thereby releasing the bioactive agent. For example, in the case of 10 kD to 30 kD PLGA, the rate of release for OP-1 is one to six weeks.

[0156] The invention also describes a method of delivering a bioactive agent at a site requiring cartilage formation comprising implanting at the defect site of a mammal a composition comprising the bioactive agent and biodegradable agent, wherein the bioactive agent is encapsulated in the biodegradable agent. Preferably, the bioactive agent is OP-1 and the biodegradable agent is PLGA.

20 EXAMPLE 1: Preparation of Tricalcium Phosphate

[0157] A slurry of lime (calcium oxide-hydroxide) is prepared and dilute phosphoric acid is added dropwise to the slurry, which is stirred constantly. The molar proportion of calcium oxide to phosphoric acid is 3:2. The product characteristics are evaluated by X-ray diffraction and adjustments are made to the proportions if required. The resultant slurry is harvested by spray drying. If the slurry is harvested by filtration, the dried cake is crushed to a fine powder of amorphous TCP. The particle size of the amorphous TCP is preferably smaller than 10  $\mu\text{m}$ .

- 68 -

EXAMPLE 2: Preparation of  $\beta$ -TCP Granule

[0158] The TCP powder was mixed with polystyrene beads (NUNC A/S-Denmark) (0-160  $\mu$ m beads). The 10 % polyvinyl pyrrolidone (PVP) granulating solution was prepared by adding PVP C-30 (Plasdone C-30, ISP technologies lot # TX 60810) in small portions in a beaker or flask of stirring water until the solution was clear. About 37 ml of 10% PVP solution was added to the TCP mixture in 5 ml increments to form a crumbly mass. As illustrated in Table 1, mixtures were prepared with different proportions of pore-forming beads and TCP.

TABLE 1

bead composition (w/w)	beads (g)	TCP (g)
12.5%	12.5	87.5
25%	12.5	37.5
37.5%	18.75	31.25
50%	23.75	23.75

[0159] The crumbly mass was passed through <500  $\mu$ m, 500-1000  $\mu$ m, or 1000-1700  $\mu$ m sieves under a vibrating motion to produce wet granules having the corresponding particle size ranges. The sieved material was dried under vacuum at 105°C for 2-3 hours.

[0160] The dried granules then underwent a burn off cycle to vaporize/carbonize the pore-forming material and were subsequently sintered at 1150°C. The temperature was raised from 39°C to 300°C over an 18 hour period, held at 300°C for 1 hour, elevated to 700°C over an 18 hour period, held at 700°C for 2 hours, and elevated to 1150°C over a 6 hour period, and held at 1150°C for 6 hours, and slow cooled to 39°C over a 6 hour period.

- 69 -

After the sintering cycle, the resultant material was confirmed by X-ray diffraction to be porous crystalline  $\beta$ -TCP.

5 [0161] The 37.5% w/w, 500-1000  $\mu$ m sintered granules were resieved and mixed with the binder, carboxy methylcellulose sodium to form a moldable putty. The putty mixtures were formed with different proportions of  $\beta$ -TCP and CMC. All combinations of  $\beta$ -TCP and CMC produced a putty having appropriate adherence properties, and did not break up in excess water. The cohesiveness of the putty was enhanced as the CMC proportion increased. The  $\beta$ -TCP/CMC 1:0.4 (w/w) putty showed the best characteristics for handling. The rheological properties of the various samples were determined.

15 EXAMPLE 3: Rat Model Bioassay for Bone Induction

[0162] This assay consists of implanting samples in subcutaneous sites in recipient rats under ether anesthesia. Male Long-Evans rats, aged 28-32 days, may be used. A vertical incision (1 cm) is made under sterile conditions in the skin over the thoracic region, and a pocket is prepared by blunt dissection. Approximately 25 mg of the test sample is implanted deep into the pocket and the incision is closed with a metallic skin clip. The day of implantation is designated as day one of the experiment. Implants are removed at varying times thereafter (i.e. 12 days, 18 days). The heterotrophic site allows for the study of bone induction without the possible ambiguities resulting from the use of orthotropic sites.

30 [0163] Bone growth is determined biochemically by calcium content of the implant. Calcium content, is proportional to the amount of bone formed in the implant. Bone formation therefore is calculated by determining the

- 70 -

calcium content of the implant in rats and is expressed as "bone forming units," where one bone forming unit represents the amount of protein that is needed for half maximal bone forming activity of the implant. Bone induction exhibited by intact demineralized rat bone matrix is considered to be the maximal bone differentiation activity for comparison purposes in this assay.

#### Cellular Events During Endochondral Bone Formation

[0164] Successful implants exhibit a controlled progression through the stages of protein-induced endochondral bone development, including: (1) transient infiltration by polymorphonuclear leukocytes; (2) mesenchymal cell migration and proliferation; (3) chondrocyte appearance; (4) cartilage matrix formation; (5) cartilage calcification; (6) vascular invasion, appearance of osteoblasts, and formation of new bone; (7) appearance of osteoclasts, bone remodeling and dissolution of the implanted matrix; and (8) hematopoietic bone marrow differentiation in the ossicles. This time course in rats may be accelerated by increasing the amounts of OP-1 added. It is possible that increasing amounts of one or more MPSFs may also accelerate this time course. The shape of the new bone conforms to the shape of the implanted matrix.

#### Histological evaluation

[0165] Histological sectioning and staining is preferred to determine the extent of osteogenesis in the implants. Implants are fixed in Bouins Solution, embedded in paraffin, and cut into 6-8  $\mu$ m sections. Staining with toluidine blue or hemotoxylin/eosin demonstrates clearly the ultimate development of

- 71 -

endochondral bone. Twelve-day implants are usually sufficient to determine whether the implants contain newly-induced bone.

#### Biological markers

5 [0166] Alkaline phosphatase (AP) activity may be used as a marker for osteogenesis. The enzyme activity may be determined spectrophotometrically after homogenization of the implant. The activity peaks at 9-10 days in vivo and thereafter slowly declines. Implants showing no bone  
10 development by histology have little or no alkaline phosphatase activity under these assay conditions. The assay is useful for quantification and obtaining an estimate of bone formation quickly after the implants are removed from the rat. Alternatively, the amount of bone  
15 formation can be determined by measuring the calcium content of the implant.

[0167] Gene expression patterns that correlate with endochondral bone or other types of tissue formation can also be monitored by quantitating mRNA levels using  
20 procedures known to those of skill in the art such as Northern Blot analysis. Such developmental gene expression markers may be used to determine progression through tissue differentiation pathways after osteogenic protein/MPSF treatments. These markers include  
25 osteoblastic-related matrix proteins such as procollagen  $\alpha_2$  (I), procollagen  $\alpha_1$ (I), procollagen  $\alpha_1$ (III), osteonectin, osteopontin, biglycan, and alkaline phosphatase for bone regeneration (see e.g., Suva et al.,  
30 J. Bone Miner. Res., 8, pp. 379-88 (1993); Benayahu et al., J. Cell. Biochem., 56, pp. 62-73 (1994)).

- 72 -

EXAMPLE 4: Sheep Model Bioassay for Bone Repair

[0168] Skeletally mature female sheep were included in the study. Three drilled defects were created in the area of the proximal metaphysis for both the left and right tibia of each animal. Defects were 6 mm in diameter and at least 10 mm deep. The defect size was consistent across all test animals. The defects were created so as to maintain the structure of the interosseous fibrofatty marrow. This marrow acts as a barrier between the implant materials and prevents interosseous mixing of the matrix materials tested. As illustrated in Table 2,  $\beta$ -TCP putty I, II, III, IV and collagen were tested in the defect sites with and without OP-1. OP-1 was either directly added to the  $\beta$ -TCP formulations or encapsulated in PLGA. Table 3 represents examples of formulations wherein the OP-1 is encapsulated in PLGA. Of the six defect sites in each animal, one defect site served as a control, which contained no test material.

[0169] A 3 to 4 inch incision was made over the proximal tibial metaphysis. The skin and underlying muscle were dissected to expose the periosteum. The periosteum was incised and maintained intact for surgical closure if possible. Three transverse holes were created in the metaphysis. The first and most superior was created approximately 2 cm below the articular surface of the tibia. The defects were created so as to form a line oriented with the long axis of the bone. Implants were spaced at 1.6 cm intervals measured center-to-center.

[0170] Materials were harvested at four and eight weeks post-treatment. Animals were euthanised with pentobarbital 75-100 mg/kg IV. The proximal tibia were taken and cut to best allow for tissue fixation. Specimens were fixed in 10% neutral buffered Formalin.

- 73 -

Specimens were cut, if feasible, so as to capture all implant sites in a single specimen. Following fixation, specimens were decalcified, embedded in plastic and sectioned in longitudinal orientation using Exackt  
 5 technique and ground to appropriate section thickness for histologic interpretation.

[0171] Radiographic assessment (Figures 9-16, 27 and 28) and histologic evaluation (Figures 1-8) were made at post-operative, four and eight weeks on all implant  
 10 sites. Anterior posterior radiographs were taken so as to best image all three defects simultaneously and view the cylindrical defects from the side. Qualitative histologic descriptions identified new bone formation, residual implant material and any evidence of pathologic  
 15 response. Images were captured for each specimen and scores presented for bone formation, acute and chronic inflammation and residual matrix.

[0172] Specimen handling and hemostatic properties were recorded at the time of implantation. Materials  
 20 ranged in form and consistency from a putty or granular form to a semi-solid cylinder.

TABLE 2

Code	Formulation	Initial pore-former percentage/Granule size
89A	$\beta$ -TCP Putty I	12.5% (w/w), 0.5-1 mm
89B	$\beta$ -TCP Putty II	25% (w/w), 0.5-1 mm
89C	$\beta$ -TCP Putty III	37.5% (w/w), 0.5-1 mm
89F	$\beta$ -TCP Putty IV	25% (w/w), 1-2 mm
48C	Collagen	Bovine type I collagen
SOB1	Lyophil 1	OP-1
SOP2	Lyophil 2	Placebo
	Reconstitution	Reconstitution medium

- 74 -

TABLE 3

Code	Formulation
formula tion 4	$\beta$ -TCP, 7% (w/w) PLGA (10 kD) with 0.3% (w/w) OP-1
5 formula tion 5	$\beta$ -TCP, 7% (w/w) PLGA (25-30 kD) with 0.3% (w/w) OP-1

Formulation Handling

[0173] Lyophil 1 and Lyophil 2 (placebo) were reconstituted by adding 2.5 ml of the reconstitution medium to one vial of the Lyophil (All components were stored frozen at 2 to 8°C until use), shaking the medium gently for 2 minutes until a homogenous (clear to cloudy) gel was formed. 0.4 ml of reconstituted Lyophil gel was added to the porous  $\beta$ -TCP matrix slowly and with care. Utilizing a thin spatula, the porous  $\beta$ -TCP matrix was mixed with the gel to form a putty-like material.

[0174] The PLGA microspheres (particle size 75-150  $\mu$ m, Alkermes, Inc.) encapsulated with 0.3% (w/w) OP-1 were mixed with the porous  $\beta$ -TCP matrix.

[0175] The putty material was immediately implanted. The implant materials were placed through the use of a folded piece of sterile paper. The paper was filled with test material and used to pour it into the defect while continuously packing material in the site. The handling properties prior to placement and in the defect site were recorded.

[0176] The  $\beta$ -TCP Putty I, II, III, IV formulations were poured as a granular dry powder. Once combined with the vehicle solution, the putties had a dry crunchy granular texture. The formulations absorbed all of the



- 75 -

Lyophil solution. The formulation was implanted with a spatula. Once in the implant site, the materials became well filled with blood.

5 [0177] The collagen formulation poured as a fluffy powder. Once mixed with a vehicle solution, it had a gritty putty texture. The formulation could be easily placed with a syringe in the implant site. The implant site became well filled with blood.

#### Histologic Results

10 [0178] Proximal tibia sections contained three defects. These defects were gross macro-cut so that all three were contained in a single section. Based on gross section observations, clinical assays, and faxitron x-rays of this section, the section was considered  
15 representative of the sample. This orientation allowed the evaluation of the periosteal reaction overlying the defects and intramedullary response to the test materials. Specimens were evaluated from 4 and 8 week explants (Figures 1-8). All three defects within a  
20 single tibial section received either the placebo or OP-1 solution. This segregation of the placebo and OP-1 implants facilitated the determination of the active or inactive biologic nature of the implant material.

#### Four-Week Evaluation for OP-1 and Placebo Implant Materials

25 [0179] At four weeks, the  $\beta$ -TCP Putty I (89A) was present in all sites (Figure 3 middle site and Figure 2 distal site). Generally, the matrix was not significantly resorbed nor was it undergoing active  
30 resorption. Sites treated with OP-1 resulted in some but not marked new bone formation (Figure 3 middle site).

- 76 -

Placebo treated sites had bone formation at the level of the cortex (Figure 2 distal site).

5 [0180] The  $\beta$ -TCP Putty II (89B) was present in all sites at 4 weeks in significant amounts (Figure 3 distal site and Figure 1 proximal site). There was no significant evidence of matrix resorption. OP-1 treated sites resulted in small amounts of new bone formation predominately at the cortical and periosteal level (Figure 3 distal site). Of the four  $\beta$ -TCP putty  
10 formulations tested,  $\beta$ -TCP putty II resulted in more inflammation than the other three formulations. Foreign body giant cells (FBGC) were reported in conjunction with this inflammation.

15 [0181]  $\beta$ -TCP Putty III (89C) was present in significant amounts in all six sites treated at 4 weeks (Figure 1 middle site and Figure 4 proximal site). OP-1 treatment did not noticeably alter residual matrix volumes. Bone formation at the cortical level was apparent in OP-1 treated specimens (Figure 4 proximal  
20 site) and less common in placebo treated sites (Figure 1 middle site). Little or no inflammation was observed in response to the  $\beta$ -TCP matrix independent of OP-1 treatment.

25 [0182]  $\beta$ -TCP Putty IV (89F) was present in significant amounts in all six sites treated at 4 weeks (Figure 1 distal site and Figure 4 middle site). OP-1 treatment had no apparent effect on residual matrix volume. OP-1 treated sites resulted in greater bone formation throughout the matrix with cortical and periosteal  
30 responses apparent (Figure 4 middle site). Little or no inflammation was observed in response to the  $\beta$ -TCP matrix independent of OP-1 treatment.

- 77 -

Eight-Week Evaluation for OP-1 and  
Placebo Treated Implant Materials

[0183] The  $\beta$ -TCP Putty I (89A) was present in all sites at 8 weeks (Figure 7 proximal site and Figure 6 distal site). The OP-1 treated implants generally showed evidence of a strong bone inductive response (Figure 7 proximal site). In two OP-1 treated sites, the  $\beta$ -TCP matrix appeared to have significantly degraded. Sites treated with OP-1 resulted in marked new bone formation at the cortical level with modest bone infiltration into the matrix within the medullary space. Placebo treated sites resulted in less bone formation at the level of the cortex (Figure 6 distal site).

[0184] The  $\beta$ -TCP Putty II (89B) was present in all sites at 8 weeks in significant amounts (Figure 5 proximal site and Figure 7 middle site). There was no significant evidence of matrix resorption. OP-1 treated sites resulted in small amounts of new bone formation predominately at the cortical and periosteal level and closure at the defect site (Figure 7 middle site). Placebo treated materials resulted in less bone formation at the cortical level and calcium particles blocking closure of the cortical defect (Figure 5 proximal site). The inflammation noted previously in response to this material was not evident.

[0185]  $\beta$ -TCP Putty III (89C) was present in significant amounts in all six sites treated at 8 weeks (Figure 5 middle site and Figure 7 distal site). OP-1 treatment did not noticeably alter residual matrix volumes. Bone formation at the cortical level and a marked periosteal response was apparent in OP-1 treated specimens (Figure 7 distal site). Little or no inflammation was observed in response to the  $\beta$ -TCP matrix independent of OP-1 treatment.

- 78 -

[0186]  $\beta$ -TCP Putty IV (89F) was present in significant amounts in all six sites treated at 8 weeks (Figure 5 distal site and Figure 8 proximal site). A few sites had less residual matrix than others. Generally, OP-1 treatment had no apparent effect on residual matrix volume. OP-1 treated sites resulted in greater bone formation throughout the matrix with an apparent cortical and periosteal response (Figure 8 proximal site). Little or no inflammation was observed in response to the  $\beta$ -TCP matrix independent of OP-1 treatment.

#### Conclusion of the Above Results

[0187] Compared to the collagen material which demonstrated acute and chronic inflammation coupled with an FBGC response, the four porous  $\beta$ -TCP formulations resulted in little or no inflammation at four and eight weeks. OP-1 treatment in the porous  $\beta$ -TCP materials consistently exhibited marked bone formation at the cortical level and a reactive periosteal response that often resulted in cortical defect closure. Although the large granular (1-2 mm)  $\beta$ -TCP putty IV formulation appeared to allow bone ingrowth deeper in the matrix, there was greater inter-granular spacing compared to that observed in small granular  $\beta$ -TCP putties.

#### Paraffin Histology Study

[0188] Tissues from the sheep model bioassay were evaluated using paraffin sections and hematoxylin and eosin stain to evaluate the effect of particle size and porosity of the implant material on bone formation in and around particles.

[0189] Tibial specimens were sectioned so as to isolate implant sites in the proximal, middle and distal sites from four animals (138, 299, 297, and 295). These

- 79 -

explants were decalcified, embedded in paraffin, sectioned and stained with hematoxylin and eosin.

[0190] Sections were viewed using light microscopy and interpreted for the effect of particle size and porosity. For specimens stratified in bone formation, the response from the cortical level was robust and deep, and the response was modest in the medullary compartment. Due to this stratification, the level extending from the endosteal cortex to a level 2-3 mm deep was evaluated.

[0191] Each of the four ceramic formulations were evaluated for bone formation in the pores and bone bridging across the particles. Bone formation in pores was assessed by counting pores that were completely isolated within a particle from the adjacent stroma. Pores that were obvious and generally round were counted. As pores were counted, a ratio was formed of those that had bone over those that did not. This is noted as the pore-fill ratio.

[0192] Pore counting was performed by scanning the field. In materials with few pores, the majority were counted as the field was scanned (Figure 25). In materials with many pores, regions were counted and a new region was viewed and then counted (Figure 26). The average of the regions or total count were presented in the ratio.

[0193] Bone bridging between particles was scored 0 to 2. A zero score was given to particles when the bone did not bridge to adjacent particles. A score of 1 was given when a couple to a few particles consistently showed bridging. A score of 2 was given when many of the particles were joined by vital bone trabeculae.

[0194] Tables 4 and 5 illustrate the pore-fill ratios and bone bridging scores for placebo and OP-1 at four weeks (Figures 17-20). Tables 6 and 7 illustrate the

- 80 -

pore-fill ratios and bone bridging scores for placebo and OP-1 at eight weeks (Figures 21-24). Bone bridging was more pronounced for  $\beta$ -TCP putties made from 37.5% (w/w) pore-forming agent and having the smaller 0.5-1 mm granule size (Tables 4-7). The pore-fill ratio was generally equivalent for the  $\beta$ -TCP putty made from 25% and 37.5% (w/w) pore-forming agents. The  $\beta$ -TCP made from 12.5% (w/w) pore-forming agent had a lower pore-fill ratio (Tables 4-7). The pore-fill ratio was consistently higher in the 89F formulation due to the larger size of the particle (1-2 mm) with more pores per particle. Compared to the small particles (0.5-1 mm), there was less bone bridging in the larger particles due to the fact that more bone was required to bridge large particles.

TABLE 4

Section	Treatment	Particle Size	Initial Pore-former %	Duration (wks)	Pore Fill Ratio	Bone Bridging
297R-D	89A	.5-1mm	12.5	4	2/10	0
297L-P	89B	.5-1mm	25	4	6/10	0
297L-M	89C	.5-1mm	37.5	4	6/7	0
297L-D	89F	1-2mm	25	4	10/10	0

Note: Section 297R-D is from the right side (R), distal (D) site of animal 297. Section 297L-P is from the left side (L), proximal site (P) of animal 297. Section 297L-M is from the left side (L), middle site (M) of animal 297. Section 297L-D is from the left side (L), distal site (D) of animal 297.

- 81 -

TABLE 5

Section	Treatment	Particle Size	Initial Pore-former %	Duration (wks)	Pore Fill Ratio	Bone Bridging
295L-M	89A	.5-1mm	12.5	4	6/11	2
295L-D	89B	.5-1mm	25	4	8/11	1
295R-P	89C	.5-1mm	37.5	4	6/8	2
295R-M	89F	1-2mm	25	4	10/10	2

Note: Section 295L-M is from the left side (L), middle (M) site of animal 295. Section 295L-D is from the left side (L), distal site (D) of animal 295. Section 295R-P is from the right side (R), proximal site (P) of animal 295. Section 295R-M is from the right side (R), middle site (M) of animal 295.

TABLE 6

Section	Treatment	Particle Size	Initial Pore-former %	Duration (wks)	Pore Fill Ratio	Bone Bridging
299R-D	89A	.5-1mm	12.5	8	4/14	1
299L-P	89B	.5-1mm	25	8	9/10	2
299L-M	89C	.5-1mm	37.5	8	18/20	2
299L-D	89F	1-2mm	25	8	9/10	1

Note: Section 299R-D is from the right side (R), distal (D) site of animal 299. Section 299L-P is from the left side (L), proximal site (P) of animal 299. Section 299L-M is from the left side (L), middle site (M) of animal 299. Section 299L-D is from the left side (L), distal site (D) of animal 299.

- 82 -

TABLE 7

Section	Treatment	Particle Size	Initial Pore-former %	Duration (wks)	Pore Fill Ratio	Bone Bridging
138L-P	89A	.5-1mm	12.5	8	10/20	1
138L-M	89B	.5-1mm	25	8	8/9	2
138L-D	89C	.5-1mm	37.5	8	10/12	2
138R-P	89F	1-2mm	25	8	9/10	1

Note: Section 138L-P is from the left side (L), proximal (P) site of animal 138. Section 138L-M is from the left side (L), middle site (M) of animal 138. Section 138L-D is from the left side (L), distal site (D) of animal 138. Section 138R-P is from the right side (R), proximal site (P) of animal 138.

#### Conclusion of Paraffin Histology Study

[0195] For  $\beta$ -TCP formulations, bone formation in pores became more apparent as the porosity increased. Bone formation in pores was less frequent in the material made from 12.5% pore-former compared to the material made from 37.5% pore-former. Although bone formation was more obvious in larger particles (1-2 mm), less bone bridging was observed in these large particles.

[0196] The collagen formulations resulted in no bone formation and a marked pathologic response. Moreover, these formulations resulted in a marked FBGCR and chronic fibroinflammatory response.

#### EXAMPLE 5: Feline Model Bioassay for Bone Repair

[0197] A femoral osteotomy defect is surgically prepared. Without further intervention, the simulated fracture defect would consistently progress to non-union. The effects of osteogenic compositions and



- 83 -

devices implanted into the created bone defects are evaluated by the following study protocol.

[0198] Briefly, the procedure is as follows: Sixteen adult cats each weighing less than 10 lbs. undergo unilateral preparation of a 1 cm bone defect in the right femur through a lateral surgical approach. In other experiments, a 2 cm bone defect may be created. The femur is immediately internally fixed by lateral placement of an 8-hole plate to preserve the exact dimensions of the defect. Four different types of materials may be implanted in the surgically created cat femoral defects: group I is a negative control group with no test material; group II is implanted with biologically active porous  $\beta$ -TCP; group III is implanted with porous  $\beta$ -TCP and an osteogenic protein; and group IV is implanted with porous  $\beta$ -TCP, an osteogenic protein and MPSF.

[0199] All animals are allowed to ambulate *ad libitum* within their cages post-operatively. All cats are injected with tetracycline (25 mg/kg subcutaneously (SQ) each week for four weeks) for bone labeling.

[0200] *In vivo* radiomorphometric studies are carried out immediately at 4, 8, 12 and 16 weeks post-operative by taking a standardized X-ray of the lightly-anesthetized animal positioned in a cushioned X-ray jig designed to consistently produce a true antero-posterior view of the femur and the osteotomy site. All X-rays are taken in exactly the same fashion and in exactly the same position on each animal. Bone repair is calculated as a function of mineralization by means of random point analysis. A final specimen radiographic study of the excised bone is taken in two planes after sacrifice.

- 84 -

[0201] Excised test and normal femurs may be immediately studied by bone densitometry, or wrapped in two layers of saline-soaked towels, placed into sealed plastic bags, and stored at -20 °C until further study.

5 Bone repair strength, load-to-failure, and work-to-failure are tested by loading to failure on a specially designed steel 4-point bending jig attached to an Instron testing machine to quantitate bone strength, stiffness, energy absorbed and deformation to failure.

10 The study of test femurs and normal femurs yields the bone strength (load) in pounds and work-to-failure in joules. Normal femurs exhibit a strength of 96 (+/- 12) pounds. Osteogenic device-implanted femur strength should be corrected for surface area at the site of

15 fracture (due to the "hourglass" shape of the bone defect repair). With this correction, the result should correlate closely with normal bone strength.

[0202] Following biomechanical testing, the bones are immediately sliced into two longitudinal sections at the

20 defect site, weighed, and the volume measured. One-half is fixed for standard calcified bone histomorphometrics with fluorescent stain incorporation evaluation, and one-half is fixed for decalcified hemotoxylin/eosin stain histology preparation.

25 [0203] Selected specimens from the bone repair site are homogenized in cold 0.15 M NaCl, 3 mM NaHCO<sub>3</sub>, pH 9.0 by a Spex freezer mill. The alkaline phosphatase activity of the supernatant and total calcium content of the acid soluble fraction of sediment are then

30 determined.

EXAMPLE 6: Rabbit Model Bioassay for Bone Repair

[0204] This assay is described in detail in Oppermann et al., U. S. Patent No. 5,354,557; see also Cook et

- 85 -

al., J. of Bone and Joint Surgery, 76-A, pp. 827-38

(1994), which are incorporated herein by reference).

Ulnar non-union defects of 1.5 cm are created in mature (less than 10 lbs) New Zealand White rabbits with epiphyseal closure documented by X-ray. The experiment may include implantation of devices into at least eight rabbits per group as follows: group I negative control implants without test material; group II implants with porous  $\beta$ -TCP; group III implants with porous  $\beta$ -TCP and an osteogenic protein; group IV implants with porous  $\beta$ -TCP, osteogenic protein and MPSF combinations. Ulnae defects are followed for the full course of the eight week study in each group of rabbits.

[0205] In another experiment, the marrow cavity of the 1.5 cm ulnar defect is packed with activated osteogenic protein in porous  $\beta$ -TCP in the presence or absence of a MPSF. The bones are allografted in an intercalary fashion. Negative control ulnae are not healed by eight weeks and reveal the classic "ivory" appearance. In distinct contrast, the osteogenic protein/MPSF-treated implants "disappear" radiographically by four weeks with the start of remineralization by six to eight weeks. These allografts heal at each end with mild proliferative bone formation by eight weeks. This type of device serves to accelerate allograft repair.

[0206] Implants treated with osteogenic protein in the presence of a MPSF may show accelerated repair, or may function at the same rate using lower concentrations of the osteogenic protein. As was described above, the rabbit model may also be used to test the efficacy of and to optimize conditions under which a particular osteogenic protein/MPSF combination can induce local bone formation.

- 86 -

EXAMPLE 7: Dog Ulnar Defect Bioassay For Bone Repair

[0207] This assay is performed essentially as described in Cook et al., Clinical Orthopaedics and Related Research, 301, pp. 302-112 (1994), which is incorporated herein by reference). Briefly, an ulnar segmental defect model is used to evaluate bone healing in 35-45 kg adult male dogs. Experimental composites comprising 500 mg of porous  $\beta$ -TCP are reconstituted with varying amounts of OP-1 in the absence or presence of increasing concentrations of one or more putative MPSFs. Any osteogenic protein may be used in place of OP-1 in this assay. Implantations at defect sites are performed with one carrier control and with the experimental series of OP-1 and OP-1/MPSF combinations being tested. Mechanical testing is performed on ulnae of animals receiving composites at 12 weeks after implantation. Radiographs of the forelimbs are obtained weekly until the animals are sacrificed at either 12 or 16 postoperative weeks. Histological sections are analyzed from the defect site and from adjacent normal bone.

[0208] The presence of one or more MPSFs may increase the rate of bone repair in dog. The presence of one or more MPSFs may also permit the use of reduced concentrations of osteogenic protein per composite to achieve similar or the same results.

EXAMPLE 8: Monkey Ulnar and Tibial Defect Bioassay For Bone Repair

[0209] This bone healing assay in African green monkeys is performed essentially as described in Cook et al., J. Bone and Joint Surgery, 77A, pp. 734-50 (1995), which is incorporated herein by reference. Briefly, a 2.0 cm osteoperiosteal defect is created in the middle of the ulnar shaft and filled with an implant comprising

- 87 -

porous  $\beta$ -TCP matrices containing OP-1 in the absence or presence of increasing concentrations of one or more putative MPSFs. Experimental composites comprising porous  $\beta$ -TCP matrices reconstituted with varying amounts of OP-1 in the absence or presence of increasing concentrations of one or more putative MPSFs are used to fill 2.0 cm osteoperiosteal defects created in the diaphysis of the tibia. Any osteogenic protein may be used in place of OP-1 in this assay. Implantations at defect sites are performed with one carrier control and with the experimental series of OP-1 and OP-1/MPSF combinations being tested. Mechanical testing is performed on ulnae and tibia of animals receiving composites. Radiographs and histological sections are analyzed from the defect sites and from adjacent normal bone as described in Cook et al.

[0210] The presence of one or more MPSFs can increase the rate of bone repair in the monkey. The presence of one or more MPSFs may also permit the use of reduced concentrations of osteogenic protein per composite to achieve similar or the same results.

EXAMPLE 9: Goat Model Fracture Healing Bioassay

[0211] This fracture healing assay in sheep is performed essentially as described in Blokhuis et al., Biomaterials, 22, pp. 725-730 (2001), which is incorporated herein by reference. A closed midshaft fracture is created in the left tibia of adult female goats with a custom-made three point bending device. The fractures are stabilized with an external fixator, which is placed at the lateral side of the tibia. Three different types of materials are implanted in the goat defects via injection: group I is a negative control group with no test material; group II is implanted with

- 88 -

the biologically active porous  $\beta$ -TCP; group III is implanted with porous  $\beta$ -TCP and an osteogenic protein; and group IV is implanted with porous  $\beta$ -TCP and an osteogenic protein encapsulated in PLGA. The test material is placed in the fractured gap. Mechanical testing (four-point non-destructive bending test) is performed on the animals receiving composites at two weeks and four weeks. After the mechanical testing, anterior, posterior, lateral, and medial slices of the fracture gap are sawn to perform radiographs and histological sections.

EXAMPLE 10: Fusion Assay of an Unstable Motor Segment of the Sheep Lumbar Spine

[0212] This assay investigates the healing of osseous and discoligamentous instability. A motor segment of the spine is a functional unit consisting of two vertebral bodies lying one above the other, and an intervertebral disc.

[0213] A trial group consists of 12 sheep. Two control groups of 12 sheep each are used. The surgical area at the inferior lumbar spine is prepared after introduction of general anesthesia and placing the animals in prone position. A skin incision of about 12 cm in length above the spinous processes of the inferior lumbar spine is made. After transsection of the subcutis and fascia, the back muscles are moved to the side.

[0214] Intubation anesthesia is applied by intramuscular injection of 1.5 ml xylazine (Rompun®). Further dosage can be administered as needed. The sedation requires placement of an intravenous indwelling catheter after puncturing an ear vein. The anesthesia is introduced through the catheter by providing 3-5 mg

- 89 -

of thiopental (Trapanal<sup>®</sup>) per killogram of body weight. After endotracheal intubation, the animals are ventilated using oxygen (30%), nitrous oxide (laughing gas) and isoflurane (Isofluran<sup>®</sup>). During the entire surgery, the analgesic fentanyl dihydrogen citrate (Fentanyl<sup>®</sup>) having a dosage 0.2-0.4 mg, is administered. At the same time, relaxation is achieved by administration of atracurium besilate (Atracurium<sup>®</sup>) at a dosage of 0.5mg/kg of body weight.

[0215] After complete exposure of the pedicles of lumbar vertebral bodies L4 to L6, a bilateral instrumentation of the pedicles L4 and L6 takes place. This is performed by using pedicle screws of 5 mm or 6 mm in diameter, depending on the diameter found in the pedicles. Subsequently, a bilateral transpedicular removal of the disc of the cranial motor segment L4/L5 is performed over the pedicle of L5 under pediculosopic control. The endplates of the affected vertebral bodies are decorticated.

[0216] Inter- and intracorporal application of test samples occurs via a transpedicular cannula in all 12 sheep of the trial group. Test samples include porous  $\beta$ -TCP, osteogenic protein or osteogenic protein encapsulated in PLGA in varying concentrations. In the first control group that consists of 12 sheep, only the porous  $\beta$ -TCP is applied. In the second control group, autologous spongiosa is administered instead of the composition of this invention.

[0217] Finally, the internal fixator is installed completely. The type of the internal fixator as well as the necessary instrumentation and surgical procedure is standardized and well known to the skilled practitioner. Drains are placed and the wound is closed using

- 90 -

absorbable suture for fascia and subcutis as well as skin staples.

5 [0218] During the entire surgical procedure, an x-ray image amplifier is available for intraoperative fluoroscopy. This facilitates exact orientation during the execution of the above steps.

10 [0219] Harvesting of the 12 sheep administered with autologous spongiosa is carried out under anesthesia as follows: the left iliac crest skin and fascia is cut by making a longitudinal incision about 8 cm long. The gluteal muscles are moved subperiostally and the cancellous bone graft is harvested from the iliac crest after an osteotomy. Bleeding control and placement of a Drain is performed upon closure of the wound in layers. The harvesting procedure is standard and known to an ordinary person skilled in the art

#### Clinical Observations

20 [0220] Daily neurologic examinations are performed to evaluate the gait of the animals as well as neurological deficits that may occur postoperatively. Operative wounds are closely examined each day. Body weights are measured preoperatively and at the time of euthanasia.

#### Radiographic Analysis

25 [0221] Before evaluation, the complete lumbar spine is freshly dissected, and the internal fixator is carefully removed. Anteroposterior and plain lateral radiographs of the operated spinal segments are obtained under consistent conditions of milliamperes, kilovolts, and seconds at 0 and 8 weeks to assist in fusion evaluation. The status of the fusion are evaluated with use of the grading system documented by Lenke et al., J.

30



- 91 -

Spinal Disord, 5, pp. 433-442 (1992), incorporated herein by reference. With this system, A indicates a big, solid trabeculated bilateral fusion mass (definitely solid); B, a big, solid unilateral fusion mass with a small contralateral fusion mass (possibly solid); C, a small, thin bilateral fusion mass with an apparent crack (probably not solid); and D, bilateral resorption of the graft or fusion mass with an obvious bilateral pseudarthrosis (definitely not solid).

[0222] Additionally, computerized tomography scans are performed to assess the fusion mass in cross sections and in saggital-plane reconstructions. For each fusion mass, approximately forty sequential computerized tomography scans are made with use of two-millimeter slice intervals and subsequent reconstruction in the saggital plane under consistent magnification and radiographic conditions.

#### Biomechanical Testing

[0223] Four specimens of each group are evaluated biomechanically. After radiographic analysis, all muscles are carefully removed while maintaining the ligamentous and bony structures. The spines are frozen at -20°C. For each of these specimens, the upper half of the upper vertebra and the lower half of the lower vertebra of the motion segment L4/L5 are embedded in polymethylmethacrylate (Technovit 3040; Heraeus Kulzer GmbH, Wehrheim/Ts, Germany). Each specimen is then fixed and tested without preload in a spine tester in a non-destructive testing mode. Alternating sequences of flexion/extension, axial right/left rotation, and right/left lateral bending moments are applied continuously at a constant rate of 1.7 degrees/second by stepper motors integrated in the gimbal of the spine

- 92 -

tester. Two precycles are applied to minimize the effect of the viscous component in the viscoelastic response, and data are collected on the third cycle. Range of motion, neutral zone, and two stiffness parameters are determined from the resulting load-deformation curves.

#### Histology/Histomorphometry

[0224] Eight specimens of each group are evaluated histologically after two, four or eight weeks postoperatively. After radiographic analysis, the spines are fixed in 10% formalin-solution. Cross sections of either specimen are obtained to evaluate bony fusion, cellular reactions, biocompatibility, and signs of cement-integration/degradation. Qualitative histologic assessment of the fusion mass at the operative site are made for the presence of giant cells, inflammatory cells, or fibrous responses where the implanted materials may have been encapsulated. In addition, the osteoid found within the trabecular fusion mass and the amount of trabecular bone are assessed. Histomorphometric variables, such as the percentage of osteoid, osteoid thickness, number of osteoblasts per millimeter bone surface, and number of osteoclasts per millimeter bone surface are determined.

#### Fluorochrome Labeling

[0225] Eight animals are subjected to intravenous application of 90 milligrams of xylenol orange per kilogram of body weight two weeks postoperatively, 10 milligrams of calcein green per kilogram of body weight four weeks postoperatively, and 25 milligrams of doxycycline hyclate yellow per kilogram of body weight six weeks postoperatively. This regimen follows the method

- 93 -

published by Rahn and Perren. See, e.g., Rahn et al., Stain Technology, 46, pp. 125-129 (1971); Rahn et al., Akt Traumatol, 10, pp. 109-115 (1980). Fluorochrome sequential analysis is then performed by Fluorescence microscopy on the specimens under UV light for qualitative and quantitative dynamic evaluation.

EXAMPLE 11: Repair of Osteochondral Defects in Dogs

[0226] A total of 12 adult male dogs are utilized. Bilateral osteochondral defects, 5.0 mm in diameter and 6 mm deep, penetrating the subchondral bone, are created in the central load bearing region of each medial femoral condyle. In 6 animals, the right defects will receive the high dose OP-1 encapsulated in PLGA. The left limb of all animals will receive the collagen matrix plus CMC to serve as a control. The remaining 6 dogs receive a low dose OP-1 encapsulated in PLGA on the right side and a control on the left side. The animals are sacrificed at 16 weeks post-implantation. At sacrifice, the distal femurs are retrieved en bloc and the defect sites are evaluated histologically and grossly based on the scheme of Moran et al., J. Bone Joint Surg. 74B, pp. 659-667 (1992), which is incorporated herein by reference.

[0227] Using standard aseptic techniques, surgery is performed under isofluorane gas anesthesia and the animals are monitored by electrocardiogram and heart rate monitors. Pre-surgical medication is administered approximately 20-30 minutes prior to anesthesia induction. The presurgical medication will consist of butorphanol tartrate (0.05 mg/kg body weight). Anesthesia is administered by intravenous injection of sodium pentothal (17.5 mg/kg body weight). Following induction, an endotracheal tube is placed and anesthesia

- 94 -

is maintained by isofluorane inhalation. Surgery is performed by making a medial parapatellar incision approximately 4 cm in length. The patella is retracted laterally to expose the femoral condyle. In the right medial condyle, a 5.0 mm diameter defect extending through the cartilage layer and penetrating the subchondral bone to a depth of 6 mm is created with a specially designed or modified 5.0 mm drill bit. After copious irrigation with saline to remove bone debris and spilled marrow cells, the appropriate concentration of OP-1 encapsulated in PLGA is carefully packed into each defect site with a blunt probe and by hand. A sufficient amount of OP-1 is placed within the defect so that it will flush with the articulating surface. While protecting the implanted material, the joint is irrigated to remove any implant not placed within the defects. The joint capsule and soft-tissues are then meticulously closed in layers with resorbable suture. The procedure is repeated on the contralateral side with placement of a control.

[0228] Butorphanol tartrate (0.05 mg/kg body weight) is administered subcutaneously as required. Animals are administered intramuscular antibiotics for four days post-surgery and routine anterior-posterior radiographs are taken immediately after surgery to insure proper surgical placement. Animals are kept in 3 x 4 feet recovery cages until the animal is able to tolerate weight bearing. Then, the animals are transferred to runs and allowed unrestricted motion.

[0229] Radiographs of the hindlimbs are obtained preoperatively, immediately postoperative, and at 16 weeks (sacrifice). The preoperative radiographs are used to assure that no pre-existing abnormalities are present and to verify skeletal maturity. Postoperative

- 95 -

radiographs are used to assess defect placement.

Sacrifice radiographs are used to assess the rate of healing and restoration of the subchondral bone and the articulating surface. Radiographs are obtained within one week of the evaluation date.

5 [0230] At the appropriate time, animals are sacrificed using an intravenous barbiturate overdose. The distal femurs are immediately harvested en bloc and stored in saline soaked towels, placed in plastic bags labeled with the animal number, right or left designation, and any other necessary identifiers. High power photographs of the defect sites are taken and carefully labeled. Prior to sacrifice venous blood is drawn for routine blood count with cell differential.

10 Soft tissues are meticulously dissected away from the defect site. The proximal end of the femur is removed. All specimens are prepared for histologic evaluation immediately after gross grading and photography. On a water cooled diamond saw each defect site is isolated.

15 [0231] The gross appearance of the defect sites and repair tissue is graded based upon the study of Moran et al., *supra*. Points are apportioned according to the presence of intra-articular adhesions, restoration of the articular surface, cartilage erosion and appearance.

20 [0232] The individual specimens are fixed by immersion in 4% paraformaldehyde solution and prepared for decalcified histologic processing. Three sections from three levels are cut from each block. Levels 1 and 3 are closest to the defect perimeter. Level 2 is located at the defect center. Three sections from each level may be stained with toluidine blue and Safranin O and fast green. Sections are graded based upon the scheme of Moran et al., *supra*. This analysis apportions points based upon the nature of the repair tissue,

25

30

- 96 -

structural characteristics, and cellular changes.

Descriptive statistics are calculated for gross and histologic parameters.

[0233] While we have described a number of  
5       embodiments of this invention, it is apparent that our  
      basic constructions may be altered to provide other  
      embodiments which utilize the products and processes of  
      this invention. Therefore, it will be appreciated that  
      the scope of this invention is to be defined by the  
10       appended claims, rather than by the specific embodiments  
      which have been presented by way of example.

- 97 -

We claim:

1. A porous  $\beta$ -TCP comprising a porous body of beta-tricalcium phosphate comprising a multiplicity of pores, wherein the pores are single separate voids having a pore diameter size of 20-500  $\mu\text{m}$ .  
5
2. A porous  $\beta$ -TCP comprising a porous body of beta-tricalcium phosphate comprising a multiplicity of pores, wherein the pores are single separate voids having a pore diameter size of 410-460  $\mu\text{m}$ .
- 10 3. A porous  $\beta$ -TCP comprising a porous body of beta-tricalcium phosphate comprising a multiplicity of pores, wherein the pores are single separate voids having a pore diameter size of 40-190  $\mu\text{m}$ .
- 15 4. A porous  $\beta$ -TCP comprising a porous body of beta-tricalcium phosphate comprising a multiplicity of pores, wherein the pores are single separate voids having a pore diameter size of 20-95  $\mu\text{m}$ .
- 20 5. A porous  $\beta$ -TCP comprising a porous body of beta-tricalcium phosphate comprising a multiplicity of pores, wherein the pores are single separate voids having a pore diameter size of 50-125  $\mu\text{m}$ .
6. The porous  $\beta$ -TCP of any one of claims 1 to 5, wherein the beta-tricalcium phosphate is sintered.
- 25 7. The porous  $\beta$ -TCP of any one of claims 1 to 5, wherein the  $\beta$ -TCP is granular and has a particle size of 0.1-2 mm.

- 98 -

8. The porous  $\beta$ -TCP of any one of claims 1 to 5, wherein the  $\beta$ -TCP is granular and has a particle size of 0.5-1.7 mm.

5 9. The porous  $\beta$ -TCP of any one of claims 1 to 5, wherein the  $\beta$ -TCP is granular and has a particle size of 1-1.7 mm.

10 10. The porous  $\beta$ -TCP of any one of claims 1 to 5, wherein the  $\beta$ -TCP is granular and has a particle size of 0.5-1.0 mm.

11. The porous  $\beta$ -TCP of any one of claims 1 to 5, wherein the total porosity is in the range of 5-80%.

12. The porous  $\beta$ -TCP of any one of claims 1 to 5, wherein the total porosity is in the range of 40-80%.

13. The porous  $\beta$ -TCP of any one of claims 1 to 5, wherein the total porosity is in the range of 65-75%.

14. The porous  $\beta$ -TCP of any one of claims 1 to 5, wherein the total porosity is 70%.

15. The porous  $\beta$ -TCP of any one of claims 1 to 5, further comprising a bioactive agent.

16. The porous  $\beta$ -TCP of claim 15, wherein the bioactive agent is a bone morphogenic protein.



- 99 -

17. The porous  $\beta$ -TCP of claim 16, wherein the bone morphogenic protein is selected from the group consisting of OP-1, OP-2, OP-3, COP-1, COP-3, COP-4, COP-5, COP-7, COP-16, BMP-2, BMP-3, BMP-3b, BMP-4, BMP-5, BMP-6, BMP-9, BMP-10, BMP-11, BMP-12, BMP-13, BMP-14, BMP-15, BMP-16, BMP-17, BMP-18, GDF-1, GDF-2, GDF-3, GDF-5, GDF-6, GDF-7, GDF-8, GDF-9, GDF-10, GDF-11, GDF-12, MP121, dorsalin-1, DPP, Vg-1, Vgr-1, 60A protein, NODAL, UNIVIN, SCREW, ADMP, NEURAL, TGF- $\beta$  and conservative amino acid sequence variants thereof having osteogenic activity.

18. The porous  $\beta$ -TCP of claim 15, wherein the bioactive agent is an osteogenic protein comprising an amino acid sequence having at least 70% homology with the C-terminal 102-106 amino acids of human OP-1.

19. The porous  $\beta$ -TCP of claim 16 further comprising a morphogenic protein stimulatory factor.

20. The porous  $\beta$ -TCP of claim 19, wherein the morphogenic protein stimulatory factor is selected from the group consisting of insulin-like growth factor I (IGF-I), estradiol, fibroblast growth factor (FGF), growth hormone (GH), growth and differentiation factor (GDF), hydrocortisone (HC), insulin, progesterone, parathyroid hormone (PTH), vitamin D, retinoic acid and IL-6.

21. The porous  $\beta$ -TCP of claim 15, wherein the bioactive agent is a nucleic acid molecule comprising a sequence encoding a bone morphogenic protein.

- 100 -

22. The porous  $\beta$ -TCP of claim 15, wherein the bioactive agent is encapsulated in a biodegradable agent.

23. The porous  $\beta$ -TCP of claim 22, wherein the  
5 biodegradable agent is selected from the group  
consisting of ethylenevinylacetate, natural and  
synthetic collagen, poly(glaxanone), poly(phosphazenes),  
polyglactin, polyglactic acid, polyaldonic acid,  
polyacrylic acids, polyalkanoates, polyorthoesters,  
10 poly(L-lactide) (PLLA), poly(D,L-lactide) (PDLLA),  
polyglycolide (PGA), poly(lactide-co-glycolide (PLGA),  
poly( $\zeta$ -caprolactone), poly(trimethylene carbonate),  
poly(p-dioxanone), poly( $\zeta$ -caprolactone-co-glycolide),  
poly(glycolide-co-trimethylene carbonate)  
15 poly(D,L-lactide-co-trimethylene carbonate),  
polyarylates, polyhydroxybutyrate (PHB), polyanhydrides,  
poly(anhydride-co-imide) and co-polymers thereof,  
polymers of amino acids, propylene-co-fumarates, a  
polymer of one or more  $\alpha$ -hydroxy carboxylic acid  
20 monomers, bioactive glass compositions, admixtures  
thereof and any derivatives and modifications thereof.

24. The porous  $\beta$ -TCP of claim 23, wherein the PLGA has a molecular weight of 5 kD to 500 kD.

25. The porous  $\beta$ -TCP of claim 23, wherein the  
25 PLGA has a molecular weight of 10 kD to 30 kD.

26. The porous  $\beta$ -TCP of claim 15, wherein the bioactive agent is an allograft or autograft.

- 101 -

27. A moldable putty composition comprising the porous  $\beta$ -TCP according to any one of claims 1 to 5 and a binder.

5                   28. The moldable putty composition of claim 27, wherein the binder is selected from the group consisting of sodium alginate, hyaluronic acid, sodium hyaluronate, gelatin, collagen, peptides, mucin, chondroitin sulfate, chitosan, poloxamer,  
10 glycosaminoglycan, polysaccharide, polyethylene glycol, methylcellulose, carboxy methylcellulose, carboxy methylcellulose sodium, carboxy methylcellulose calcium, hydroxypropyl methylcellulose, hydroxybutyl methylcellulose, hydroxyethyl methylcellulose,  
15 hydroxyethylcellulose, methylhydroxyethyl cellulose, hydroxyethyl cellulose, polylactic acid, polyglycolic acid, co-polymers of polylactic acid and polyglycolic acid, polyhydroxybutyric acid, polymalic acid, polyglutamic acid, polylactone, mannitol, white  
20 petrolatum, mannitol/dextran combinations, mannitol/white petrolatum combinations, sesame oil, fibrin glue and admixtures thereof.

25                   29. The moldable putty composition of claim 28, wherein the fibrin glue is a mixture of human fibrinogen and thrombin.

30. The moldable putty composition of claim 27 further comprising a bioactive agent.

31. A kit comprising:  
a) the porous  $\beta$ -TCP of any one of claims  
30 1 to 5; and  
b) a bioactive agent.

- 102 -

32. The kit of claim 31, wherein the bioactive agent is a bone morphogenic protein.

33. The kit of claim 32, wherein the bone morphogenic protein is selected from the group  
5 consisting of OP-1, OP-2, OP-3, COP-1, COP-3, COP-4, COP-5, COP-7, COP-16, BMP-2, BMP-3, BMP-3b, BMP-4, BMP-5, BMP-6, BMP-9, BMP-10, BMP-11, BMP-12, BMP-13, BMP-14, BMP-15, BMP-16, BMP-17, BMP-18, GDF-1, GDF-2, GDF-3, GDF-5, GDF-6, GDF-7, GDF-8, GDF-9, GDF-10,  
10 GDF-11, GDF-12, MP121, dorsalin-1, DPP, Vg-1, Vgr-1, 60A protein, NODAL, UNIVIN, SCREW, ADMP, NEURAL, TGF- $\beta$  and conservative amino acid sequence variants thereof having osteogenic activity.

34. The kit of claim 31, wherein the  
15 bioactive agent is an osteogenic protein comprising an amino acid sequence having at least 70% homology with the C-terminal 102-106 amino acids of human OP-1.

35. The kit of claim 32 further comprising a morphogenic protein stimulatory factor.

20 36. The kit of claim 35, wherein the morphogenic protein stimulatory factor is selected from the group consisting of insulin-like growth factor I (IGF-I), estradiol, fibroblast growth factor (FGF), growth hormone (GH), growth and differentiation factor (GDF), hydrocortisone (HC), insulin, progesterone,  
25 parathyroid hormone (PTH), vitamin D, retinoic acid and IL-6.

- 103 -

37. A kit comprising:

- a) the porous  $\beta$ -TCP of any one of claims 1 to 5; and
- b) a binder.

5                   38. The kit of claim 37, wherein the binder ,  
is selected from the group consisting of sodium  
alginate, hyaluronic acid, sodium hyaluronate, gelatin,  
collagen, peptides, mucin, chondroitin sulfate,  
10                   chitosan, poloxamer, glycosaminoglycan, polysaccharide,  
polyethylene glycol, methylcellulose, carboxy  
methylcellulose, carboxy methylcellulose sodium, carboxy  
methylcellulose calcium, hydroxypropyl methylcellulose,  
hydroxybutyl methylcellulose, hydroxyethyl  
15                   methylcellulose, hydroxyethylcellulose,  
methylhydroxyethyl cellulose, hydroxyethyl cellulose,  
polylactic acid, polyglycolic acid, co-polymers of  
polylactic acid and polyglycolic acid,  
polyhydroxybutyric acid, polymalic acid, polyglutamic  
20                   acid, polylactone, mannitol, white petrolatum,  
mannitol/dextran combinations, mannitol/white petrolatum  
combinations, sesame oil, fibrin glue and admixtures  
thereof.

25                   39. The kit of claim 38, wherein the fibrin  
glue is a mixture of human fibrinogen and thrombin.

40. An implantable prosthetic device  
comprising:

- a) a prosthetic implant having a  
surface region implantable adjacent to a target tissue;  
30                   and
- b) the porous  $\beta$ -TCP of any one of  
claims 1 to 5 disposed on the surface region.

- 104 -

41. The prosthetic device of claim 40 further comprising a bioactive agent dispersed in the porous  $\beta$ -TCP.

5 42. The prosthetic device of claim 41, wherein the bioactive agent is a bone morphogenic protein.

43. The prosthetic device of claim 42, wherein the bone morphogenic protein is selected from the group consisting of OP-1, OP-2, OP-3, COP-1, COP-3, COP-4, COP-5, COP-7, COP-16, BMP-2, BMP-3, BMP-3b, BMP-4, BMP-5, BMP-6, BMP-9, BMP-10, BMP-11, BMP-12, BMP-13, BMP-14, BMP-15, BMP-16, BMP-17, BMP-18, GDF-1, GDF-2, GDF-3, GDF-5, GDF-6, GDF-7, GDF-8, GDF-9, GDF-10, GDF-11, GDF-12, MP121, dorsalin-1, DPP, Vg-1, Vgr-1, 60A protein, NODAL, UNIVIN, SCREW, ADMP, NEURAL, TGF- $\beta$  and conservative amino acid sequence variants thereof having osteogenic activity.

44. The prosthetic device of claim 41, wherein the bioactive agent is an osteogenic protein comprising an amino acid sequence having at least 70% homology with the C-terminal 102-106 amino acids of human OP-1.

45. The prosthetic device of claim 42 further comprising a morphogenic protein stimulatory factor.

25 46. The prosthetic device of claim 45, wherein the morphogenic protein stimulatory factor is selected from the group consisting of insulin-like growth factor I (IGF-I), estradiol, fibroblast growth factor (FGF), growth hormone (GH), growth and

- 105 -

differentiation factor (GDF), hydrocortisone (HC), insulin, progesterone, parathyroid hormone (PTH), vitamin D, retinoic acid and IL-6.

5           47. The prosthetic device of claim 41, wherein the bioactive agent is a nucleic acid molecule comprising a sequence encoding a bone morphogenic protein.

10           48. The prosthetic device of claim 41, wherein the bioactive agent is encapsulated in a biodegradable agent.

15           49. The prosthetic device of claim 48, wherein the biodegradable agent is selected from the group consisting of ethylenevinylacetate, natural and synthetic collagen, poly(glaxanone), poly(phosphazenes), polyglactin, polyglactic acid, polyaldonic acid, polyacrylic acids, polyalkanoates, polyorthoesters, poly(L-lactide) (PLLA), poly(D,L-lactide) (PDLA), polyglycolide (PGA), poly(lactide-co-glycolide (PLGA), poly( $\zeta$ -caprolactone), poly(trimethylene carbonate), 20 poly(p-dioxanone), poly( $\zeta$ -caprolactone-co-glycolide), poly(glycolide-co-trimethylene carbonate) poly(D,L-lactide-co-trimethylene carbonate), polyarylates, polyhydroxybutyrate (PHB), polyanhydrides, poly(anhydride-co-imide) and co-polymers thereof, 25 polymers of amino acids, propylene-co-fumarates, a polymer of one or more  $\alpha$ -hydroxy carboxylic acid monomers, bioactive glass compositions, admixtures thereof and any derivatives and modifications thereof.

- 106 -

50. The prosthetic device of claim 49,  
wherein the PLGA has a molecular weight of 5 kD to 500 kD.

51. The prosthetic device of claim 49,  
wherein the PLGA has a molecular weight of 10 kD to 30  
5 kD.

52. The prosthetic device of claim 40,  
wherein the device is selected from the group consisting  
of a hip device, a fusion cage and a maxillofacial  
device.

10 53. The prosthetic device of claim 40 further  
comprising a binder.

54. The prosthetic device of claim 53,  
wherein the binder is selected from the group consisting  
of sodium alginate, hyaluronic acid, sodium hyaluronate,  
15 gelatin, collagen, peptides, mucin, chondroitin  
sulfate, chitosan, poloxamer, glycosaminoglycan,  
polysaccharide, polyethylene glycol, methylcellulose,  
carboxy methylcellulose, carboxy methylcellulose sodium,  
carboxy methylcellulose calcium, hydroxypropyl  
20 methylcellulose, hydroxybutyl methylcellulose,  
hydroxyethyl methylcellulose, hydroxyethylcellulose,  
methylhydroxyethyl cellulose, hydroxyethyl cellulose,  
polylactic acid, polyglycolic acid, co-polymers of  
polylactic acid and polyglycolic acid,  
25 polyhydroxybutyric acid, polymalic acid, polyglutamic  
acid, polylactone, mannitol, white petrolatum,  
mannitol/dextran combinations, mannitol/white petrolatum  
combinations, sesame oil, fibrin glue and admixtures  
thereof.



- 107 -

55. The prosthetic device of claim 54, wherein the fibrin glue is a mixture of human fibrinogen and thrombin.

5 56. A method of producing a porous  $\beta$ -TCP granule comprising:

(a) blending a TCP powder with a pore-forming agent;

(b) adding a granulating solution to form a crumbly mass;

10 (c) passing the crumbly mass through a sieve to form granules; and

(d) sintering the granules to form porous  $\beta$ -TCP.

15 57. A method of producing a porous  $\beta$ -TCP granule comprising:

(a) blending a TCP powder with a pore-forming agent, wherein the proportion of pore-forming agent is 37.5% by weight;

20 (b) adding a granulating solution to form a crumbly mass;

(c) passing the crumbly mass through a sieve to form granules; and

(d) sintering the granules to form porous  $\beta$ -TCP.

25 58. A method of producing a porous  $\beta$ -TCP granule comprising:

(a) blending a TCP powder with a pore-forming agent;

30 (b) adding a granulating solution to form a crumbly mass;

- 108 -

(c) passing the crumbly mass through a sieve to form granules, wherein the sieve is in the size range of 500-1000  $\mu\text{m}$  or 1000-1700  $\mu\text{m}$ ; and

5 (d) sintering the granules to form porous  $\beta$ -TCP.

59. A method of producing a porous  $\beta$ -TCP granule comprising:

(a) blending a TCP powder with a pore-forming agent;

10 (b) adding a granulating solution to form a crumbly mass;

(c) passing the crumbly mass through a sieve to form granules;

15 (d) vaporizing the granules at 700-800°C; and

(e) sintering the granules to form porous  $\beta$ -TCP.

60. A method of producing a porous  $\beta$ -TCP granule comprising:

20 (a) blending a TCP powder with a pore-forming agent;

(b) adding a granulating solution to form a crumbly mass;

25 (c) passing the crumbly mass through a sieve to form granules; and

(d) sintering the granules at 1000-1200°C and followed by a slow cooling step to form porous  $\beta$ -TCP.

30 61. The method of any one of claims 56 to 60, wherein the pore-forming agent is selected from the group consisting of prepolymers of polyacrylates,

- 109 -

polymethacrylates, polymethyl methacrylate, copolymers of methyl acrylate and methyl methacrylate, polystyrene, polyethylene glycol, crystalline cellulose, fibrous cellulose, polyurethanes, polyethylenes, nylon resins and acrylic resins.

62. The method of any one of claims 56 to 60, wherein the granulating solution comprises a compound selected from the group consisting of polyvinyl pyrrolidone, starch, gelatin, polyvinyl alcohol, polyethylene oxide, hydroxyethyl cellulose, polyvinyl butyral and cellulose acetate butyrate.

63. The method of any one of claims 56 to 60, wherein the porous  $\beta$ -TCP is resieved after formation.

64. A composition comprising tricalcium phosphate powder and a pore-forming agent, wherein the pore-forming agent has a diameter of 20-500  $\mu\text{m}$ .

65. A composition comprising tricalcium phosphate powder and a pore-forming agent, wherein the pore-forming agent has a diameter of 410-460  $\mu\text{m}$ .

66. A composition comprising tricalcium phosphate powder and a pore-forming agent, wherein the pore-forming agent has a diameter of 40-190  $\mu\text{m}$ .

67. A composition comprising tricalcium phosphate powder and a pore-forming agent, wherein the pore-forming agent has a diameter of 20-95  $\mu\text{m}$ .

- 110 -

68. A composition comprising tricalcium phosphate powder and a pore-forming agent, wherein the pore-forming agent has a diameter of 50-125  $\mu\text{m}$ .

5 69. The composition of any one of claims 64 to 68, wherein the proportion of pore-forming agent is 30-40% by weight.

10 70. A method of inducing bone formation in a mammal comprising the step of implanting in the defect site of said mammal a composition comprising the porous  $\beta$ -TCP according to any one of claims 1 to 5.

71. The method of claim 70, wherein the composition further comprises a bioactive agent.

72. The method of claim 71, wherein the bioactive agent is a bone morphogenic protein.

15 73. The method of claim 70, wherein the composition further comprises a binder.

20 74. A method of delivering a bioactive agent at a site requiring bone formation comprising implanting at the defect site of a mammal a composition comprising the porous  $\beta$ -TCP of any one of claims 1 to 5 and a bioactive agent.

75. The method of claim 74, wherein the bioactive agent is a bone morphogenic protein.

25 76. The method of claim 74, wherein the bioactive agent is encapsulated in a biodegradable agent.

- 111 -

77. The method of claim 76, wherein the delivery of the bioactive agent is sustained released.

5 78. The method of claim 74, wherein the bioactive agent is a nucleic acid molecule comprising a sequence encoding a bone morphogenic protein.

10 79. A method of delivering a bioactive agent at a site requiring cartilage formation comprising implanting at the defect site of a mammal a composition comprising a bioactive agent and a biodegradable agent having a particle size of 20-500  $\mu\text{m}$ , wherein the bioactive agent is encapsulated in the biodegradable agent.

15 80. A composition comprising a bioactive agent encapsulated in a biodegradable agent, wherein the biodegradable agent has a particle size of 20-500  $\mu\text{m}$ .

81. A composition comprising a bioactive agent encapsulated in a biodegradable agent, wherein the biodegradable agent has a particle size of 20-140  $\mu\text{m}$ .

20 82. A composition comprising a bioactive agent encapsulated in a biodegradable agent, wherein the biodegradable agent has a particle size of 75-140  $\mu\text{m}$ .

25 83. The composition of any one of claims 80 to 82, wherein the biodegradable agent is selected from the group consisting of ethylenevinylacetate, natural and synthetic collagen, poly(glaxanone), poly(phosphazenes), polyglactin, polyglactic acid, polyaldonic acid, polyacrylic acids, polyalkanoates, polyorthoesters, poly(L-lactide) (PLLA),

- 112 -

poly(D,L-lactide) (PDLA), polyglycolide (PGA),  
poly(lactide-co-glycolide (PLGA), poly( $\zeta$ -caprolactone),  
poly(trimethylene carbonate), poly(p-dioxanone),  
poly( $\zeta$ -caprolactone-co-glycolide),  
5 poly(glycolide-co-trimethylene carbonate)  
poly(D,L-lactide-co-trimethylene carbonate),  
polyarylates, polyhydroxybutyrate (PHB), polyanhydrides,  
poly(anhydride-co-imide) and co-polymers thereof,  
polymers of amino acids, propylene-co-fumarates, a  
10 polymer of one or more  $\alpha$ -hydroxy carboxylic acid  
monomers, bioactive glass compositions, admixtures  
thereof and any derivatives and modifications thereof.

84. The composition of claim 83, wherein the  
PLGA has a molecular weight of 5 kD to 500 kD.

15 85. The composition of claim 83, wherein the  
PLGA has a molecular weight of 10 kD to 30 kD.



Figure 1




Figure 2



3/28



Figure 3

4/28



Figure 4

5/28



Figure 5




Figure 6

7/28




Figure 7

8/28




Figure 8

9/28



Figure 9

10/28




Figure 10



11/28



Figure 11

12/28




Figure 12

13/28



Figure 13

14/28



Figure 14

15/28



Figure 15

16/28




Figure 16

17/25



Figure 17

18/25



Figure 18



19/28




Figure 19

20/28



Figure 20

21/28



Figure 21

22/28




Figure 22



Figure 23

24/28



Figure 24

25/28



Figure 25



Figure 26



27/28




Figure 27

28/28




Figure 28

## SEQUENCE LISTING

<110> Stryker Corporation et al.  
 <120> POROUS BETA-TRICALCIUM PHOSPHATE GRANULES AND METHODS  
 FOR PRODUCING SAME  
 <130> STK-8 PCT  
 <140> Not Assigned  
 <141> Concurrently herewith  
 <150> 09/798,518,  
 09/960,789  
 <160> 2001-03-02,  
 2001-09-21  
 <160> 10  
 <170> PatentIn Ver. 2.1  
 <210> 1  
 <211> 431  
 <212> PRT  
 <213> Homo sapiens  
 <400> 1  
 Met His Val Arg Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala  
 1 5 10 15  
 Leu Trp Ala Pro Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser  
 20 25 30  
 Leu Asp Asn Glu Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser  
 35 40 45  
 Gln Glu Arg Arg Glu Met Gln Arg Glu Ile Leu Ser Ile Leu Gly Leu  
 50 55 60  
 Pro His Arg Pro Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro  
 65 70 75 80  
 Met Phe Met Leu Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Gly Gly  
 85 90 95  
 Gly Pro Gly Gly Gln Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser  
 100 105 110  
 Thr Gln Gly Pro Pro Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr  
 115 120 125  
 Asp Ala Asp Met Val Met Ser Phe Val Asn Leu Val Glu His Asp Lys  
 130 135 140  
 Glu Phe Phe His Pro Arg Tyr His His Arg Glu Phe Arg Phe Asp Leu  
 145 150 155 160  
 Ser Lys Ile Pro Glu Gly Glu Ala Val Thr Ala Ala Glu Phe Arg Ile  
 165 170 175

Tyr Lys Asp Tyr Ile Arg Glu Arg Phe Asp Asn Glu Thr Phe Arg Ile  
                   180                                  185                                  190  
 Ser Val Tyr Gln Val Leu Gln Glu His Leu Gly Arg Glu Ser Asp Leu  
                   195                                  200                                  205  
 Phe Leu Leu Asp Ser Arg Thr Leu Trp Ala Ser Glu Glu Gly Trp Leu  
                   210                                  215                                  220  
 Val Phe Asp Ile Thr Ala Thr Ser Asn His Trp Val Val Asn Pro Arg  
                   225                                  230                                  235                                  240  
 His Asn Leu Gly Leu Gln Leu Ser Val Glu Thr Leu Asp Gly Gln Ser  
                                   245                                  250                                  255  
 Ile Asn Pro Lys Leu Ala Gly Leu Ile Gly Arg His Gly Pro Gln Asn  
                                   260                                  265                                  270  
 Lys Gln Pro Phe Met Val Ala Phe Phe Lys Ala Thr Glu Val His Phe  
                   275                                  280                                  285  
 Arg Ser Ile Arg Ser Thr Gly Ser Lys Gln Arg Ser Gln Asn Arg Ser  
                   290                                  295                                  300  
 Lys Thr Pro Lys Asn Gln Glu Ala Leu Arg Met Ala Asn Val Ala Glu  
                   305                                  310                                  315                                  320  
 Asn Ser Ser Ser Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr  
                                   325                                  330                                  335  
 Val Ser Phe Arg Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu  
                                   340                                  345                                  350  
 Gly Tyr Ala Ala Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn  
                   355                                  360                                  365  
 Ser Tyr Met Asn Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His  
                   370                                  375                                  380  
 Phe Ile Asn Pro Glu Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln  
                   385                                  390                                  395                                  400  
 Leu Asn Ala Ile Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile  
                                   405                                  410                                  415  
 Leu Lys Lys Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys His  
                   420                                  425                                  430

&lt;210&gt; 2

&lt;211&gt; 96

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

 <223> Description of Artificial Sequence: Synthetic  
           amino acid sequence

&lt;400&gt; 2

Leu Tyr Val Asp Phe Ser Asp Val Gly Trp Asp Asp Trp Ile Val Ala  
   1                                  5                                  10                                  15

Pro Pro Gly Tyr Gln Ala Phe Tyr Cys His Gly Glu Cys Pro Phe Pro  
                   20                  25                  30  
 Leu Ala Asp His Phe Asn Ser Thr Asn His Ala Val Val Gln Thr Leu  
                   35                  40                  45  
 Val Asn Ser Val Asn Ser Lys Ile Pro Lys Ala Cys Cys Val Pro Thr  
                   50                  55                  60  
 Glu Leu Ser Ala Ile Ser Met Leu Tyr Leu Asp Glu Asn Glu Lys Val  
                   65                  70                  75                  80  
 Val Leu Lys Tyr Asn Gln Glu Met Val Val Glu Gly Cys Gly Cys Arg  
                   85                  90                  95

<210> 3  
 <211> 96  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
           amino acid sequence

<400> 3  
 Leu Tyr Val Asp Phe Ser Asp Val Gly Trp Asn Asp Trp Ile Val Ala  
   1                  5                  10                  15  
 Pro Pro Gly Tyr His Ala Phe Tyr Cys His Gly Glu Cys Pro Phe Pro  
                   20                  25                  30  
 Leu Ala Asp His Leu Asn Ser Thr Asn His Ala Val Val Gln Thr Leu  
                   35                  40                  45  
 Val Asn Ser Val Asn Ser Lys Ile Pro Lys Ala Cys Cys Val Pro Thr  
                   50                  55                  60  
 Glu Leu Ser Ala Ile Ser Met Leu Tyr Leu Asp Glu Asn Glu Lys Val  
                   65                  70                  75                  80  
 Val Leu Lys Tyr Asn Gln Glu Met Val Val Glu Gly Cys Gly Cys Arg  
                   85                  90                  95

<210> 4  
 <211> 102  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: OPX

<220>  
 <221> MOD\_RES  
 <222> (1)..(102)  
 <223> Xaa is independently selected from a group of one  
           or more specified amino acids as defined in the  
           specification

<400> 4

Cys Xaa Xaa His Glu Leu Tyr Val Xaa Phe Xaa Asp Leu Gly Trp Xaa  
 1 5 10 15  
 Asp Trp Xaa Ile Ala Pro Xaa Gly Tyr Xaa Ala Tyr Tyr Cys Glu Gly  
 20 25 30  
 Glu Cys Xaa Phe Pro Leu Xaa Ser Xaa Met Asn Ala Thr Asn His Ala  
 35 40 45  
 Ile Xaa Gln Xaa Leu Val His Xaa Xaa Xaa Pro Xaa Xaa Val Pro Lys  
 50 55 60  
 Xaa Cys Cys Ala Pro Thr Xaa Leu Xaa Ala Xaa Ser Val Leu Tyr Xaa  
 65 70 75 80  
 Asp Xaa Ser Xaa Asn Val Xaa Leu Xaa Lys Xaa Arg Asn Met Val Val  
 85 90 95  
 Xaa Ala Cys Gly Cys His  
 100

<210> 5  
 <211> 97  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Generic  
 Sequence 7

<220>  
 <221> MOD\_RES  
 <222> (1)..(97)  
 <223> Xaa is independently selected from a group of one  
 or more specified amino acids as defined in the  
 specification

<400> 5  
 Leu Xaa Xaa Xaa Phe Xaa Xaa Xaa Gly Trp Xaa Xaa Xaa Xaa Xaa  
 1 5 10 15  
 Pro Xaa Xaa Xaa Xaa Ala Xaa Tyr Cys Xaa Gly Xaa Cys Xaa Xaa Pro  
 20 25 30  
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Asn His Ala Xaa Xaa Xaa Xaa  
 35 40 45  
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Cys Xaa Pro  
 50 55 60  
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Xaa Xaa Xaa Xaa Xaa Xaa  
 65 70 75 80  
 Val Xaa Leu Xaa Xaa Xaa Xaa Xaa Met Xaa Val Xaa Xaa Cys Xaa Cys  
 85 90 95  
 Xaa

<210> 6

<211> 102  
 <212> PRT  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Generic  
 Sequence 8

<220>

<221> MOD\_RES

<222> (1)..(102)

<223> Xaa is independently selected from a group of one  
 or more specified amino acids as defined in the  
 specification

<400> 6

Cys Xaa Xaa Xaa Xaa Leu Xaa Xaa Xaa Phe Xaa Xaa Xaa Gly Trp Xaa  
 1 5 10 15

Xaa Xaa Xaa Xaa Xaa Pro Xaa Xaa Xaa Xaa Ala Xaa Tyr Cys Xaa Gly  
 20 25 30

Xaa Cys Xaa Xaa Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Asn His Ala  
 35 40 45

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
 50 55 60

Xaa Cys Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Xaa Xaa  
 65 70 75 80

Xaa Xaa Xaa Xaa Xaa Val Xaa Leu Xaa Xaa Xaa Xaa Xaa Met Xaa Val  
 85 90 95

Xaa Xaa Cys Xaa Cys Xaa  
 100

<210> 7  
 <211> 97  
 <212> PRT  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Generic  
 Sequence 9

<220>

<221> MOD\_RES

<222> (1)..(97)

<223> Xaa is independently selected from a group of one  
 or more specified amino acids as defined in the  
 specification

<400> 7

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
 1 5 10 15

Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Gly Xaa Cys Xaa Xaa Xaa  
 20 25 30

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
           35                          40                          45  
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Pro  
           50                          55                          60  
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
           65                          70                          75                          80  
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Cys  
                           85                          90                          95  
 Xaa

<210> 8  
 <211> 102  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Generic  
           Sequence 10

<220>  
 <221> MOD\_RES  
 <222> (1)..(102)  
 <223> Xaa is independently selected from a group of one  
           or more specified amino acids as defined in the  
           specification

<400> 8

Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
   1                          5                          10                          15  
 Xaa Xaa Xaa Xaa Xaa Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Gly  
           20                          25                          30  
 Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
           35                          40                          45  
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
           50                          55                          60  
 Xaa Xaa Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
           65                          70                          75                          80  
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
           85                          90                          95  
 Xaa Xaa Cys Xaa Cys Xaa  
           100

<210> 9  
 <211> 5  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Consensus



## sequence

&lt;220&gt;

&lt;221&gt; MOD\_RES

&lt;222&gt; (2)..(5)

<223> Xaa is independently selected from a group of one  
or more specified amino acids as defined in the  
specification

&lt;400&gt; 9

Cys Xaa Xaa Xaa Xaa

1 5

&lt;210&gt; 10

&lt;211&gt; 1822

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (49)..(1341)

&lt;400&gt; 10

ggtgcggggcc cggagcccgg agcccgggta gcgcgtagag ccggcgcg atg cac gtg 57  
Met His Val  
1

cgc tca ctg cga gct gcg gcg ccg cac agc ttc gtg gcg ctc tgg gca 105  
Arg Ser Leu Arg Ala Ala Pro His Ser Phe Val Ala Leu Trp Ala  
5 10 15

ccc ctg ttc ctg ctg cgc tcc gcc ctg gcc gac ttc agc ctg gac aac 153  
Pro Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser Leu Asp Asn  
20 25 30 35

gag gtg cac tcg agc ttc atc cac cgg cgc ctc cgc agc cag gag cgg 201  
Glu Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser Gln Glu Arg  
40 45 50

cgg gag atg cag cgc gag atc ctc tcc att ttg ggc ttg ccc cac cgc 249  
Arg Glu Met Gln Arg Glu Ile Leu Ser Ile Leu Gly Leu Pro His Arg  
55 60 65

ccg cgc ccg cac ctc cag ggc aag cac aac tcg gca ccc atg ttc atg 297  
Pro Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro Met Phe Met  
70 75 80

ctg gac ctg tac aac gcc atg gcg gtg gag gag ggc ggc ggg ccc ggc 345  
Leu Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Gly Gly Gly Pro Gly  
85 90 95

ggc cag ggc ttc tcc tac ccc tac aag gcc gtc ttc agt acc cag ggc 393  
Gly Gln Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser Thr Gln Gly  
100 105 110 115

ccc cct ctg gcc agc ctg caa gat agc cat ttc ctc acc gac gcc gac 441  
Pro Pro Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr Asp Ala Asp  
120 125 130

atg gtc atg agc ttc gtc aac ctc gtg gaa cat gac aag gaa ttc ttc	489
Met Val Met Ser Phe Val Asn Leu Val Glu His Asp Lys Glu Phe Phe	
135 140 145	
cac cca cgc tac cac cat cga gag ttc cgg ttt gat ctt tcc aag atc	537
His Pro Arg Tyr His His Arg Glu Phe Arg Phe Asp Leu Ser Lys Ile	
150 155 160	
cca gaa ggg gaa gct gtc acg gca gcc gaa ttc cgg atc tac aag gac	585
Pro Glu Gly Glu Ala Val Thr Ala Ala Glu Phe Arg Ile Tyr Lys Asp	
165 170 175	
tac atc cgg gaa cgc ttc gac aat gag acg ttc cgg atc agc gtt tat	633
Tyr Ile Arg Glu Arg Phe Asp Asn Glu Thr Phe Arg Ile Ser Val Tyr	
180 185 190 195	
cag gtg ctc cag gag cac ttg ggc agg gaa tcg gat ctc ttc ctg ctc	681
Gln Val Leu Gln Glu His Leu Gly Arg Glu Ser Asp Leu Phe Leu Leu	
200 205 210	
gac agc cgt acc ctc tgg gcc tcg gag gag ggc tgg ctg gtg ttt gac	729
Asp Ser Arg Thr Leu Trp Ala Ser Glu Glu Gly Trp Leu Val Phe Asp	
215 220 225	
atc aca gcc acc agc aac cac tgg gtg gtc aat ccg cgg cac aac ctg	777
Ile Thr Ala Thr Ser Asn His Trp Val Val Asn Pro Arg His Asn Leu	
230 235 240	
ggc ctg cag ctc tcg gtg gag acg ctg gat ggg cag agc atc aac ccc	825
Gly Leu Gln Leu Ser Val Glu Thr Leu Asp Gly Gln Ser Ile Asn Pro	
245 250 255	
aag ttg gcg ggc ctg att ggg cgg cac ggg ccc cag aac aag cag ccc	873
Lys Leu Ala Gly Leu Ile Gly Arg His Gly Pro Gln Asn Lys Gln Pro	
260 265 270 275	
ttc atg gtg gct ttc ttc aag gcc acg gag gtc cac ttc cgc agc atc	921
Phe Met Val Ala Phe Phe Lys Ala Thr Glu Val His Phe Arg Ser Ile	
280 285 290	
cgg tcc acg ggg agc aaa cag cgc agc cag aac cgc tcc aag acg ccc	969
Arg Ser Thr Gly Ser Lys Gln Arg Ser Gln Asn Arg Ser Lys Thr Pro	
295 300 305	
aag aac cag gaa gcc ctg cgg atg gcc aac gtg gca gag aac agc agc	1017
Lys Asn Gln Glu Ala Leu Arg Met Ala Asn Val Ala Glu Asn Ser Ser	
310 315 320	
agc gac cag agg cag gcc tgt aag aag cac gag ctg tat gtc agc ttc	1065
Ser Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr Val Ser Phe	
325 330 335	
cga gac ctg ggc tgg cag gac tgg atc atc gcg cct gaa ggc tac gcc	1113
Arg Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu Gly Tyr Ala	
340 345 350 355	
gcc tac tac tgt gag ggg gag tgt gcc ttc cct ctg aac tcc tac atg	1161
Ala Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn Ser Tyr Met	
360 365 370	

```

aac gcc acc aac cac gcc atc gtg cag acg ctg gtc cac ttc atc aac 1209
Asn Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His Phe Ile Asn
      375                      380                      385

ccg gaa acg gtg ccc aag ccc tgc tgt gcg ccc acg cag ctc aat gcc 1257
Pro Glu Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln Leu Asn Ala
      390                      395                      400

atc tcc gtc ctc tac ttc gat gac agc tcc aac gtc atc ctg aag aaa 1305
Ile Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile Leu Lys Lys
      405                      410                      415

tac aga aac atg gtg gtc cgg gcc tgt ggc tgc cac tagctcctcc 1351
Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys His
      420                      425                      430

gagaattcag accctttggg gccaaagtttt tctggatcct ccattgctcg ccttggccag 1411

gaaccagcag accaactgcc ttttgtgaga ccttcccctc cctatcccca actttaaagg 1471

tgtgagagta ttaggaaaca tgagcagcat atggcttttg atcagttttt cagtggcagc 1531

atccaatgaa caagatccta caagctgtgc aggcaaaacc tagcaggaaa aaaaaacaac 1591

gcataaagaa aaatggccgg gccagggtcat tggctgggaa gtctcagcca tgcacggact 1651

cgtttccaga ggtaattatg agcgcctacc agccaggcca cccagccgtg ggaggaaggg 1711

ggcgtggcaa ggggtgggca cattgggtgtc tgtgcgaaaag gaaaattgac ccggaagtgc 1771

ctgtaataaa tgtcacaata aaacgaatga atgaaaaaaaa aaaaaaaaaa a 1822

```